

Journal of Coastal Life Medicine

journal homepage: www.jclmm.com



Document heading

doi:10.12980/JCLM.2.2014C1177

© 2014 by the Journal of Coastal Life Medicine. All rights reserved.

Traditional ethno–botanical uses of medicinal plants from coastal areas of Pakistan

Muhammad Qasim¹, Zainul Abideen¹, Muhammad Yousuf Adnan¹, Raziuddin Ansari¹, Bilquees Gul^{1*}, Muhammad Ajmal Khan^{1,2}

¹Institute of Sustainable Halophyte Utilization, University of Karachi–75270, Karachi, Pakistan

²Qatar Shell Professorial Chair of Sustainable Development, Department of International Affairs, College of Arts and Sciences, Qatar University, PO Box 2713, Doha, Qatar

PEER REVIEW

Peer reviewer

Prof. (Emer.) Dr. Münir Öztürk, Ege University, Botany Department, Izmir, Turkey.

Tel: 0090–535–3098104

E-mail: munirozturk@gmail.com

Comments

This investigation has enlightened the fact that documentation of therapeutic uses of herbal remedies with new phyto–medicinal claim is very important. This study will form the bases for detailed studies on the bioactive compounds in these plants by pharmacologists. It may result in the discovery of novel compounds for health services
Details on Page 28

ABSTRACT

Objective: To document the traditional uses of wild plants as medicine by the villagers along the coastal highway from Karachi to Uthal.

Methods: Information presented in this research was gathered from the local people using an integrated approach of floral collections, discussions with the elderly people and traditional medicinal practitioners using semi–structured questionnaire.

Results: Ethno–medicinal surveys indicated the medicinal importance of 54 plant species from 27 families in the targeted area. Majority of the plants (54%) from this coastal plant diversity were xerophytes followed by halophytes/xero–halophytes (40%) and glycophytes (6%). The most important uses included gastrointestinal diseases, pain killer, arthritis, skin and sexual disorders, asthma and expectorant. The above–ground parts of plants *i.e.* leaf, stem and fruit/seed as decoction are used most commonly to cure 23 ailments but root was also used in some cases.

Conclusions: This study helps in documenting therapeutic uses of herbal remedies with new phyto–medicinal claim and it is hoped that it will lead to detailed chemical and pharmacological evaluations. This may also lead to a discovery of novel bioactive compounds for food and pharmaceutical industries.

KEYWORDS

Arabian Sea, Halophytes, Herbal medicine, Karachi, Salt tolerant, Uthal

1. Introduction

Awareness about the therapeutic properties of plants is existing since few thousand years ago^[1]. However, use of plants to cure diseases and relieve physical sufferings dates back to the earlier times of mankind’s history probably starting from the first moment when a human being or their animal got ill. It is also reported that expeditions, undertaken by those interested in the cultural attributes of primitive societies, obtained valuable information regarding the uses

of plants for medicine^[2]. The use of plants as medicine by traditional people laid basis for the evolution of modern medicine^[3]. The American Shaman Inc. is an example of a company that has focused on traditional medical systems in their drug discovery programs.

Herbal therapy has come of age and today, medicinal plants play a significant role in human health care globally^[4]. In USA about 25% of all prescriptions dispensed in public pharmacies contain drugs extracted from higher plants and about 64% of the total global population remains

*Corresponding author: Bilquees Gul, Institute of Sustainable Halophyte Utilization, University of Karachi–75270, Karachi, Pakistan.

E-mail: bilqueesgul@uok.edu.pk; gul_bilquees@yahoo.com

Tel: +9221–34820253

Fax: +9221–34820258

Foundation Project: Supported by Pakistan Academy of Sciences (Grant no: 5–9/PAS, 2010).

Article history:

Received 5 Jan 2014

Received in revised form 12 Jan, 2nd revised form 15 Jan, 3rd revised form 19 Jan 2014

Accepted 23 Jan 2014

Available online 28 Jan 2014

dependent on traditional medicine for their healthcare needs^[3]. In India alone, approximately 7 500 medicinal plants are traditionally used against various ailments while Korea, Japan, China, and Malaysia together with few other southeast Asian countries are the other leading in the world in the use of herbal medicines^[5,6]. Flora of Pakistan reported to consist of about 5000 species and less than 10% of them are used either as herbal medicine or used in about 350 classical formulations to treat various diseases^[7]. Pakistan represented a classical case study for investigating the role of traditional medicine for multiple healthcare settings. It is a predominantly rural country with 67% population living in remote areas taking advantage of local flora in pluralistic means^[8]. Rural settlements along coastal belt of Pakistan (Sindh and Balochistan provinces) presented a classical example of having rich history of using herbal medicine. The area is a native home of herbal plants renowned for providing primary health care in traditional system. A variety of wild plants have been collected and sold in the local market by native communities. However, very limited scientific knowledge is available on the potential herbs, which can be cultivated and utilized for different purposes. This is however, practiced without any technical knowledge and hence there is a need to adopt a scientific approach for preserving this heritage.

Phyto-geographical studies conducted in the coastal areas of Sindh and Balochistan revealed that most of the plants found here were either halophytes or xero-halophytes, due to arid/semi-arid environment and saline nature of the soil^[9,10]. To cope with abiotic stresses under such conditions, plants are known to synthesize secondary metabolites which can be a rich source of medicinally important compounds such as flavonoids, terpenoids, tannins, alkaloids *etc*^[11]. Furthermore, these plants are source of food, fodder, medicine, biofuels, edible oil *etc.* in this region and they are proving their potentials to be cultivated as cash crops in many parts of the world^[10-16]. Despite these credentials, these plants had been relatively under explored for exploiting their potential.

Therefore, this study has been undertaken to gather available information on the medicinal uses of these plants by rural communities along coastal areas to treat different diseases and to provide scientific basis for their effectiveness.

2. Materials and methods

The traditional ethno-medicinal information was collected

from Karachi (Sindh) to Uthal (Balochistan) during 2008 to 2011 to document the medicinal uses of wild plants. The area is characterized by dry and saline land; where brackish underground water is available and the dominant vegetation was either halophytic or xerophytic.

Information was collected from traditional healers, herbalists and rural dwellers of the villages and interviews were conducted by using semi-structured questionnaire^[17]. For this purpose, priority was given to elders because they have better knowledge than young generation. During these interviews and transect walks, information about particular plants was collected for their part used *i.e.* above or below ground, method of preparation, details of administration, the dosage *etc.* Plant materials were collected from their natural populations during organized tours accompanied by herbalists or any knowledgeable rural dwellers. Plants initially identified by their local names but later proper identification was done at the herbarium, University of Karachi. Identified plants were deposited in the herbarium as voucher specimens for later access.

3. Results

3.1. The plants and their medical application

The present study have shown that traditional medicine is still playing a significant role in meeting fundamental healthcare needs of inhabitants around coastal area of Pakistan. The survey gathered information on 54 plant species reported by the informants for their medicinal uses (Table 1 and Table 2). The reported species were distributed among 27 botanical families (Table 1). Amaranthaceae (6 species) was most represented family, followed by Mimosaceae (4 species) while Asclepidaceae, Asteraceae, Caesalpinaceae, Papilionaceae and Poaceae comprises 3 species (each) and rest of the families included two and one member.

The most important uses related to gastrointestinal diseases (12% species), followed by pain killer (9% species), arthritis (8% species), skin diseases and sexual disorders (7% species), asthma and expectorant (6% species). The rest were used to treat internal and external problems such as alexipharmic, eye infections, hepatitis, jaundice, malaria, tuberculosis, UGT disorders and ulceration (Figure 1).

Analysis of the data revealed that a single species is used for multiple health problems like *Citrullus colocynthis* and *Eucalyptus glandulosa* were applied extensively against wide range of ailments (Table 2). They were used against

six and five different ailments respectively (Figure 1). The rest of the medicinal plants had one to four medicinal uses (Figure 1).

Stomach disorder was the most common health care problem reported in this study (Figure 1). About 12 different plant species were used to treat different digestive disorders. In this regard the most frequently used plants were *Acacia nilotica*, *Capparis decidua*, *Chenopodium album*, *Desmostachya bipinnata*, *Grewia tenax*, *Indigofera*

oblongifolia and *Prosopis cineraria*. The mode of preparation of the remedy was decoction in water whereas in case of fruits it was consumed directly.

Utilization of medicinal plants as a remedy to cure pain is quite common only second to stomach disorder. About 9 species have been reported to utilize in this category in which *Calotropis procera*, *Fagonia indica*, *Haloxylon stocksii*, *Solanum surattense*, *Tribulus terrestris* and *Zizyphus nummularia* were some common examples.

Table 1

List of medicinal plants with respect to their family, local name, habit, life form and plant type.

Name of specie	Family	Local name	Habit	Life Form	Plant Type
<i>Abutilon indicum</i> (L.) Sweet, Hort. Brit.	Malvaceae	Khangi	Shrub	Perennial	Xerophyte
<i>Acacia nilotica</i> (L.) Delile	Mimosaceae	Babool/Desi Keekar	Tree	Perennial	Xerophyte
<i>Acacia senegal</i> L.	Mimosaceae	Kumbat	Tree	Perennial	Xerophyte
<i>Achyranthes aspera</i> L.	Amaranthaceae	Kandii	Herb	Annual	Xerophyte
<i>Aeluropus lagopoides</i> (L.) Trin. ex Thw.	Poaceae	Pooji Chabbar	Grass	Perennial	Halophyte
<i>Aerva javanica</i> (Brum. f.) Juss. Ex J.A. Schultes var. <i>bovei</i> Webb.	Amaranthaceae	Boo	Herb	Perennial	Xerophyte
<i>Aerva javanica</i> (Brum. f.) Juss. ex J.A. Schultes var. <i>javanica</i>	Amaranthaceae	Boo	Herb	Perennial	Xerophyte
<i>Aizoon canariense</i> L.	Aizoaceae	Welaiti battar/Dotak	Herb	Annual	Xerohlophite
<i>Alhaji maurorum</i> Medic.	Papilionaceae	Kandaira	Shrub	Perennial	Halophyte
<i>Ammi visnaga</i> (L.) Lam.	Apiaceae	Spairkai	Herb	Annual	Xerophyte
<i>Artemisia scoparia</i> Waldst. & Kit.	Asteraceae	Jaanh/Jaukay	Herb	Perennial	Xerophyte
<i>Arthrocnemum indicum</i> (Willd.) Moq.	Amaranthaceae	Namak zor	Herb	Perennial	Halophyte
<i>Azadirachta indica</i> Adr. Juss.	Meliaceae	Neem	Tree	Perennial	Xerophyte
<i>Caesalpinia bonduc</i> (L.) Roxb.	Caesalpinaceae	Karbat	Tree	Perennial	Xerophyte
<i>Calotropis procera</i> (Ait.) Ait.	Asclepidaceae	Aak	Shrub to tree	Perennial	Xerophyte
<i>Capparis decidua</i> Forssk.	Capparidaceae	Kareer	Shrub to tree	Perennial	Xerophyte
<i>Cassia holosericea</i> Fresen	Caesalpinaceae	Goora wall	Shrub	Perennial	Xerophyte
<i>Chenopodium album</i> L.	Amaranthaceae	Lullar	Herb	Annual	Halophyte
<i>Citrullus colocynthis</i> (L.) Schrad	Cucurbitaceae	Timmay or Too	Herb	Perennial	Xerophyte
<i>Cleome viscosa</i> L.	Capparidaceae	Kinni Buti	Herb	Annual	Xerophyte
<i>Convolvulus glomeratus</i> Choisy	Convolvulaceae	Rakewall	Herb	Perennial	Xerophyte
<i>Corchorus depressus</i> (L.) Stocks	Tiliaceae	Mundairy	Herb	Perennial	Xerophyte
<i>Cressa cretica</i> L.	Convolvulaceae	Bukkan	Herb	Annual	Halophyte
<i>Cymbopogon juvarancusa</i> (Jones) Schult.	Poaceae	Nadak	Grass	Perennial	Halophyte
<i>Cyperus rotundus</i> L.	Cyperaceae	Dodi Ghass	Grass	Annual	Halophyte
<i>Desmostachya bipinnata</i> (L.) Stapf	Poaceae	Drub	Grass	Perennial	Halophyte
<i>Enicostema hyssopifolium</i> (Willd) Verdoo	Gentianaceae	Mamraycho	Herb	Annual	Xerohalophyte
<i>Eucalyptus camaldulensis</i> L.	Myrtaceae	Baam or Safaida	Tree	Perennial	Halophyte
<i>Euphorbia caducifolia</i> Haines	Euphorbiaceae	Thowwar	Shrub	Perennial	Xerophyte
<i>Euphorbia hirta</i> L.	Euphorbiaceae	Kippo	Herb	Annual	Xerophyte
<i>Fagonia indica</i> ssp. <i>schweinfurthia</i> Hadidi	Zygophyllaceae	Kandayra	Shrublet	Perennial	Xerophyte
<i>Glossonema varians</i> (Stocks) Hook. f.	Asclepiadaceae	Munga	Herb	Annual	Xerophyte
<i>Grewia tenax</i> (Forsk.) Fiori	Tiliaceae	Gangi	Shrub	Perennial	Xerophyte
<i>Haloxylon stocksii</i> (Boiss.) Benth. & Hook.	Amaranthaceae	Khaar	Shrub	Perennial	Xerohalophyte
<i>Indigofera oblongifolia</i> Forsk.	Papilionaceae	Chiel	Shrub	Perennial	Xerohalophyte
<i>Indigofera cordifolia</i> Heyne ex Roth	Papilionaceae	Mangoli	Herb	Annual	Xerohalophyte
<i>Inula grantioides</i> Boiss	Asteraceae	Colmeer	Herb	Perennial	Xerophyte
<i>Launaea resedifolia</i> (L.) Kuntze	Asteraceae	Badtar	Herb	Annual	Halophyte
<i>Leptadenia pyrotechnica</i> (Forssk.) Dcne.	Asclepidaceae	Kheep	Shrub	Perennial	Xerophyte
<i>Mangifera indica</i> L.	Anacardiaceae	Aam	Tree	Perennial	Glycophyte
<i>Parkinsonia aculeata</i> L.	Caesalpinaceae	Welaiti keekar	Tree	Perennial	Xerohalophyte
<i>Portulaca oleracea</i> L.	Portulacaceae	Kulfa	Herb	Annual	Xerophyte
<i>Prosopis cineraria</i> (L.) Druce	Mimosaceae	Kandu	Shrub to tree	Perennial	Xerophyte
<i>Prosopis juliflora</i> (Swartz) DC.	Mimosaceae	Davi	Shrub to tree	Perennial	Xerohalophyte
<i>Punica granatum</i> L.	Punicaceae	Darroo/Annar	Shrub to tree	Perennial	Glycophyte
<i>Salvadora oleoides</i> Dne.	Salvadoraceae	Jaar	Shrub to tree	Perennial	Xerophyte
<i>Salvadora persica</i> L.	Salvadoraceae	Jaar	Shrub to tree	Perennial	Xerophyte
<i>Solanum surattense</i> Burm. f.	Solanaceae	Kandarii	Herb	Perennial	Xerohalophyte
<i>Syzygium cumini</i> (L.) Skeels	Myrtaceae	Jamman	Tree	Perennial	Glycophyte
<i>Thespesia populneoides</i> (Roxb.) Kostel.	Malvaceae	Peepal	Tree	Perennial	Halophyte
<i>Trianthema portulacastrum</i> L.	Aizoaceae	Itsit	Herb	Annual	Xerohalophyte
<i>Tribulus terrestris</i> L.	Zygophyllaceae	Kandawaro	Herb	Annual	Xerophyte
<i>Withania somnifera</i> (L.) Dunal	Solanaceae	Ashgand	Shrub	Perennial	Xerohalophyte
<i>Zizyphus nummularia</i> (Burm. f.) Wight and Arn.	Rhamnaceae	Jungli Bayri	Shrub to tree	Perennial	Xerophyte

Table 2

Ethno-botanical uses of medicinal plants from coastal areas and vicinity.

Name of specie	Part used	Used against	Preparation	Dosage	R A	Reported uses	References
<i>Abutilon indicum</i>	L	Painful urination and painful menses	Decoction	Twice a day	O	Antimicrobial	[22]
<i>Acacia nilotica</i>	Le/F	Dysentery, diarrhea, vomiting, acidity	Decoction	1 spoon 3 times a day until cure	O	Asthma, diarrhea, demulcent	[11]
<i>Acacia senegal</i>	L	Burns, wounds, inflammation	Paste	Once a day	E	Antimicrobial	[23]
<i>Achyranthes aspera</i>	L	Diarrhea, dysentery, vomiting, abdominal cramp, ulcer	Decoction	3 times a day for a week	O	Antibacterial, Antifungal	[24]
<i>Aeluropus lagopoides</i>	WP	Wound healing, pain killer	Apply paste	Until cure	E	Not reported	
<i>Aerva javanica</i> var <i>bovei</i>	I	Alexipharmic	Take inflorescence, warm it and place it on the site of bite	Use until cure	E	Diuretic activity	[25]
		Asthma, headache, rheumatism	Make pillow of inflorescence	Use until cure	E	Antioxidant	[26]
<i>Aerva javanica</i> var <i>javanica</i>	I	Alexipharmic	Take inflorescence, warm it and place it on the site of bite	Use until cure	E	Wounds, Jaundice, Diabetes, cough	[11]
		Asthma, headache, rheumatism	Make pillow of inflorescence	Use until cure	E		
<i>Aizoon canariensis</i>	WP	Jaundice, hepatitis	Take 250–300 g plant material in 1/2 liter of rapeseed oil and massage on chest	3–4 times a day	E	Not reported	
<i>Alhagi maurorum</i>	WP	Laxative, expectorant	Decoction	Twice a day	O	Antiinflammatory, Antipyretic, Antioxidant	[27]
<i>Ammi visnaga</i>	F	Heart problems, expectorant	Decoction	Twice a day	O	Antioxidant	[28]
<i>Artemisia scoparia</i>	WP	Burns	Smoke is used against Burns	Twice a day	E	Antifungal, Insecticidal	[29,30]
<i>Arthrocnemum indicum</i>	WP	Alexipharmic	Ash of plant	Until cure	E	Alexipharmic	[31,11]
<i>Azadiracta indica</i>	L	Headache	Paste	Use until cure	E	Antioxidant, Antiviral	[32]
	F	Hemorrhoids	Eat raw fruits	Use until cure	O	Antibacterial	
<i>Caesalpinia bonduc</i>	S	Resettlement of disturbed joints and bones specially after trauma, arthritis	Decoction	1 cup daily	O	Antimicrobial	[33]
<i>Calotropis procera</i>	L	Pain killer specially for chest and ribs	Add few drops of oil on leaf then warm it and bandage over affected area	3 times a day	E	Antioxidant, Antibacterial	[34]
	R	Malarial fever	Decoction of root with black pepper	2–3 times a day	O	Tooth and stomach aches	[11]
<i>Capparis decidua</i>	F	Laxative, stomach worms	Eat unripe fruits without seeds	Few fruits once in a day	O	Antifungal, Carminative, aphrodisiac, ulcer, cough, asthma	[11]
	S	Toothache, gum infection	Powder of young stem	Twice a day	ATG		[36]
	L	Hepatitis, jaundice, ulcers,	Decoction	Twice a day	O	Cathartic activity	[37]
<i>Cassia holosericea</i>	L	Laxative	Decoction	Once a day in fasting	O		
	Se	Laxative	Decoction	Once a day in fasting	O		
<i>Chenopodium album</i>	WP	Dysentery, diarrhea, headache	Decoction	Twice a day	O	Antimicrobial Anthelmintic	[38]
	F	T.B, asthma, whooping cough	Fruit filled with Ajwain seeds and left for a week in dark then use the Ajwain seeds	1 spoon daily	O	Antioxidant, Anti-diabetic, Antiinflammatory	[39,40]
<i>Citrullus colocynthis</i>	F	Whole body swelling/elephantiasis	Massage with paste of fruits	Once a day	E	Toothache, Constipation, Piles	[11]
	F	Eye irritation, inflammation	Take an eye pick and embedded in fruit then apply on eyes	Twice a day	E	Leucorrhea, Asthma	
<i>Cleome viscosa</i>	R	Toothache, gum inflammation, infection	Miswak of roots	2–3 times a day	ATG		
	L	Infertility	Infusion	Once a day	O	Anthelmintic	[41]
<i>Convolvulus glomeratus</i>	WP	Laxative	Decoction with two drops of <i>Calotropis procera</i> latex	Two spoons only	O	Not reported	
<i>Corchorus depressus</i>	WP	Spermatorrhoea, painful urination, sexual disorders, impotenc	Infusion	One glass daily for one week	O	Antipyretic Analgesic	[42]
<i>Cressa cretica</i>	WP	Expectorant, asthma	Decoction	Twice a day	O	Antiinflammatory, Antioxidant, Antiviral	[43]
<i>Cymbopogon jwarancusa</i>	WP	Productive cough, chest itching, Bronchitis, pharyngitis	Decoction	One cup twice a day	O	Antipyretic activity	[44]
<i>Cyperus rotundus</i>	R	Boils, blisters	Powder	Twice a day	E	Antioxidant, Antiinflammatory	[45]
<i>Desmostachya bipinnata</i>	R	Dysentery	Decoction	3 times a day	O	Antimicrobial, Carbuncle	[11,46]
<i>Enicostema hysopifolium</i>	WP	Malarial and Dengue fever	Decoction	1 glass 3 times a day	O	Antimicrobial	[47]
		Pain killer	Paste	Massage	E	Hypoglycemic activity	[48]
<i>Eucalyptus camaldulensis</i>	L/S	Fever, cough, sneezing, spasmodic, malarial shivering	Decoction of leaves and paste of stem	2 times a day	O/N	Antimicrobial	[49]
	L	Pain killer specially for bones and joints	Paste	Massage twice a day	E	Larvicidal	[50]
<i>Euphorbia caducifolia</i>	La	Pain killer specially for bones	Milky Latex use for massage	Twice a day	E	Milk clotting activity	[11,51]
	S	Alexipharmic	Cut stem, warm it and place on the site of bite	Use until cure	E	Increase male sexual vigor	
<i>Euphorbia hirta</i>	WP	Asthma	Decoction	Taken frequently	O	Antibacterial	[52]
<i>Fagonia indica</i>	WP	Severe itching	Decoction	Take bath	O	Antitumor activity	[53]
	WP	Pain killer	Decoction in milk	One glass twice a day	O	Analgesic Antimicrobial	[54]
<i>Glossonema varians</i>	L	Painful urination	Infusion	One glass for 10 days	O	Not reported	
<i>Grewia tenax</i>	F/Se	Stomachache specially in stomach ulcer	Raw	10–20 g daily	O	Antibacterial	[55]
<i>Haloxylon stocksii</i>	WP	Arthritis, resettlement of disturb joints	Decoction used for massage	Massage	E	Paraplegic limbs	[11]
<i>Indigofera oblongifolia</i>	R	Dysentery, diarrhea, vomiting, acidity, ulcer	Decoction	3 times a day	O	Antibacterial Antifungal	[56]

Table 2 continued,

Ethno-botanical uses of medicinal plants from coastal areas and vicinity.

Name of specie	Part used	Used against	Preparation	Dosage	RA	Reported uses	Reference
	L	Boils, blisters	Paste of leaves	Apply paste and bandage	E	Antiinflammatory, Cytotoxic activity	[57]
<i>Indigofera cordifolia</i>	WP	Dysentery, diarrhea, vomiting	Decoction	2 spoons 3 times a day	O	Wound healing activity, Antiviral	[58]
<i>Inula granioides</i>	WP	Prolong fever, rheumatism	Boil in water and bath	Take bath daily	B	Not reported	
		Abortifacient	Decoction	Only 2 glass	O		
<i>Launea residifolia</i>	L	Jaundice, leucorrhoea	Decoction	1 spoon 3 times a day	O	Antibacterial	[59]
<i>Leptadenia pyrotechnica</i>	S/WP	For UGT disorders, Dysmenorrhea, Spermatorrhia, impotency	Decoction	Half cup twice a day for a week	O	Antioxidant Antimicrobial	[60]
<i>Mangifera indica</i>	B	Dysentery, diarrhea, vomiting	Infusion	1–2 times daily	O	Antibacterial	[61]
<i>Parkinsonia aculeata</i>	L	Arthritis, rheumatism, pain killer	Infusion	1 glass twice a day for 2 weeks	O	Antidiabetic Antimalaria	[62]
<i>Portulaca oleracea</i>	L F/Se	Blood purification Toothache	Decoction Paste	One cup daily twice a day	O ATC	Antioxidant, Antibacterial	
<i>Prosopis cineraria</i>	L	Diarrhea, dysentery, vomiting, abdominal cramp	Raw and decoction	Few leaves or 1 spoon 3 times a day	O	Hypoglycemic, Antioxidative Antimicrobial	[64]
<i>Prosopis juliflora</i>	S	Boils, blisters, skin infections	Burn young stem from one side, it release extract on the other side. Apply extract on infected area.	3–4 times	E	Immunomodulating, Antidermatophytic, Antibacterial	[65]
	L	Nail infection	Crush leaves and add few ml of sunflower oil and bandage over infected area	Once	E	Cytotoxic activity, Antiinflammatory, Antimicrobial	[66]
<i>Punica granatum</i>	FS	Dysentery, vomiting, food poisoning	Infusion	Twice a day	O	Antimicrobial	[67]
	F	Heart attack, cardiovascular problems	Fruit	Fistful twice a day	O	Antioxidant	[68]
<i>Salvadora oleoides</i>	S	Toothache, gum infection	Use as Miswak or stem powder	3 times a day	ATC	Antiinflammatory	[69]
<i>Salvadora persica</i>	S	Toothache, gum infection	Use as Miswak or stem powder	3 times a day	ATC	Antiinflammatory	[70]
<i>Solanum surattense</i>	F	Arthritis, resettlement of disturb joints	Paste with egg white	Massage	E	Antibacterial	[71]
<i>Syzygium cumini</i>	L	Dysentery, diarrhea, bloody stools	Infusion	3 times a day	O	Antidiarrhoeal	[72]
	L	Skin infections, wounds	Paste	Twice a day	E		
<i>Thespesia populnea</i>	L	Hepatitis, jaundice, ulcers	Decoction	Twice a day	O	Antimicrobial	[73]
<i>Trianthema portulacastrum</i>	L	Asthma	Decoction	Half cup Twice a day	O	Antioxidant Antiinflammatory	[74]
<i>Tribulus terrestris</i>	F	Arthritis, backache	Make a sweet dish (Halwa)	3 times a day	O	Antimicrobial	[75]
	F	Spermatorrhoea, impotence	Powder of dry fruits with sugar and black pepper	Twice a day	O	Aphrodisiac activity, Skin diseases	[76,77]
<i>Withania somnifera</i>	R	Spermatorrhia, debility	Root powder with honey and long pepper	One spoon daily	O	Antimicrobial, Body vigor	[78,79]
<i>Zizyphus nummularia</i>	F	Arthritis, backache	Make a sweet dish (Halwa)	3 times a day	O	Antibacterial, Antifungal	[80]

Key: B–Bark; F–Fruit; FS–Fruit Skin; I–Inflorescence; L–Leaves; La–Latex ; L–Legume; R–Root; S–Stem; Se–Seed; WP–Whole plant; E–Externally; O–Orally; O/N–Oral and Nasal Breath; ATC; Apply on Teeth and Gums; RA–route of Administration.




Figure 1. Distribution of medicinal plants with respect to a: diseases, b: number of uses, c: plant type, d: plant part used, e: mode of preparation, f: route of administration.

Consequently, a number of preparations were offered to treat infertility, vaginal infections, child bearing, delivery, painful urination and painful menses, impotency and debility, where on the other hand, some other healers provided women with prescriptions for abortion and contraception. Child health received considerable attention from healers and a number of herbs were prescribed for treating baby indigestion, cramps, jaundice, dehydration, constipation, eye infections and flu (Table 2).

3.2. Plant type, life form and habit of the medicinal plants

Distribution of medicinal plants with respect to their plant type showed (Figure 1) that about 54% of the reported taxa belonged to xerophytes and 40% from halophytes (including halophytes and xerohalophytes) and few of them were glycophytes (6%). The predominant habit (Figure 1) reported in this study was herbs (43%) followed by shrubs (31%) trees (19%) and grasses (7%). Most of the reported plants were perennials (72%) followed by annuals (28%) since rainfall is a rare and unpredictable event in arid and semi-arid areas.

3.3. Plant part used and mode of preparation

All parts of the various plants were used for the preparation

of different herbal formulations; however, whole plant (24%), fruit (23%) and leaves (21%) were the most frequently plant parts used. Sometimes, the stem (10%), root (9%), seeds (5%) and inflorescence (3%) were also used (Figure 1). The legume, gum or raisin, bark and latex were also reported in some cases.

Fresh or dry plant samples both were used to prepare herbal formulations. The most commonly used method was decoction (40%) followed by direct consumption of raw plant parts (23%) then paste of plant material (13%) and in some cases maceration technique as infusion in water (11%) was also used. Furthermore powder of a plant material, preparation of a sweet dish (Halwa), ashes and smoke of the plants were also used in herbal medication (Figure 1).

3.4. Route of administration and dosage

The formulation and administration of herbal remedy depends upon the type of therapeutic condition. Remedies prescribed for curing skin diseases and traumas were applied externally, whereas medication of internal problems exclusively involves oral administration. However, some of skin diseases and certain traumas also require oral administration. Therefore, most of the drugs prescribed by traditional healers are administered orally. About 62% of local remedies were taken orally, followed by 30% which are of external use (applied topically on skin) and 8% are applied exclusively on teeth and gums (Figure 1).

4. Discussion

This research summarized ethno-botanical uses of medicinal plants distributed around the coastal areas of Pakistan (from Karachi to Uthal). The area is characterized by high salinity, temperature and light intensity with low rain fall. Plants growing in such environments are mostly halophytes and xerophytes which are well adapted to survive under extreme conditions^[10,11]. They evolve multiple mechanisms including synthesis of diverse group of distinctive compounds which help them to grow in conditions where other plants failed to survive^[9,13]. These chemicals when utilized can serve as potent medicine against different ailments. Their range of active ingredients utilized traditionally to combat deadly ailments and these plants serve as a successful home remedy for deprived people living along the coast. A Field-based, ethnographic observation of coastal areas revealed that most of the rural people do not have access to health care professionals. They used a wide range of locally available plant species against multiple diseases^[10,11]. The common uses were against

stomach related problems, body pain, arthritis, skin and sexual disorders, asthma *etc.* The most widespread problem was digestive disorder which reflected the real condition of rural settlements. Lack of basic amenities along with poor sanitation, unhygienic conditions and consumption of contaminated water are some basic reasons for digestive system disorders which may be the root causes for most of the internal diseases^[18].

Present investigation also revealed that the local people make good use of a wide range of plants from their surroundings. All the 54 plants had one or more medicinal uses where some of them used for 6 different conditions in which every plant part used as exclusive medicinal remedy^[7]. To attain a positive response of local medication, one should select the appropriate part of a plant that contains the active ingredients and properly prepare the formulation^[4]. Among all these preparations, a standardized decoction method was widely used. Decoction is a most traditional method in which the plant material is boiled in water to extract all essential compounds and other medicinal substances to get fast relief. However, lack of precision was observed to determine the dose to be taken and it was considered a drawback of herbal medication. In most of the cases the medicinal formulations were taken orally and it was also observed that most of the documented preparations were drawn from a single plant where mixtures were used rarely.

It is also a matter of deep concern that the transmission of traditional knowledge is decreasing through several generations because of no proper way of documenting this heritage except to specific individuals interested in preserving their traditions^[8]. On the other hand the local medicinal plants are under threat due to anthropogenic and developmental activities such as urbanization; road buildings, growth of mega-structures *etc.* are examples of direct assault on nature resulting in the loss of local flora^[1,3]. Therefore, it is strongly recommended that the proper documentation of traditional knowledge, utilization of natural resources and conservation of unique floral wealth should be done in a scientific way to prevent their losses.

Present study concluded that communities along the coast line of Pakistan inherit a rich traditional knowledge and provided novel information pertaining to local medicinal flora. This study provides detailed account of therapeutic uses of herbal remedies including 54 species with new phyto-medicinal claims, which would be a significant ethno-pharmaceutical contribution. These plants need immediate consideration for carrying out detailed chemical and pharmacological evaluations. Such investigations may lead to the discovery of novel bioactive compounds that will help to assess the efficacy of herbal remedies.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgements

Thanks are owed to the herbalist, local practitioners and rural dwellers for their cooperation, sharing their knowledge and guiding in fields. The research work is supported by Pakistan Academy of Sciences (Grant no: 5–9/PAS, 2010). The first two authors are recipient of scholarship from Higher Education Commission of Pakistan under their Indigenous PhD. fellowship program.

Comments

Background

Halophytes are plants which grow all over the world, in different climatic regions on areas with high soil salinity and play an important role in the ecological balance as well as preservation of habitats. Their economic potential has emerged very slowly and is attracting the attention of many researchers to understand their role in the development of human resources.

Research frontiers

Modern medical facilities in rural areas are generally lacking in the villages and people have to rely on herbal drugs. An attempt has been made here to document the traditional uses of wild plants as medicine by the villagers along the coastal highway from Karachi to Uthal in Pakistan.

Related reports

As a general practice of the ethnobotanical knowledge collection information here too has been gathered from the locals living in the area, floral collections undertaken, and a semi-structured questionnaire filled up during the discussions with the elderly people and traditional medicinal practitioners.

Innovations and breakthroughs

In all 54 plant species from 27 families have been collected from the coast, majority being xerophytes followed by halophytes/xero-halophytes. The report is first attempt from this unique habitat. It is well documented with tables and figures.

Applications

Most important applications are gastrointestinal diseases, pain killer, arthritis, skin and sexual disorders, asthma and expectorant. The leaf, stem and fruit/seed as decoction are most commonly used to cure 23 ailments.

Peer review

This investigation has enlightened the fact that documentation of therapeutic uses of herbal remedies with new phyto-medicinal claim is very important. This study will form the bases for detailed studies on the bioactive compounds in these plants by pharmacologists. It may result in the discovery of novel compounds for health services

References

- [1] Jones FA. Herbs—useful plants. Their role in history and today. *Eur J Gastroenterol Hepatol* 1996; **8**: 1227–1231.
- [2] Lal SD, Yadav BK. Folk medicines of Kurukshetra District (Haryana), Indian. *Eco Bot* 1983; **37**: 299–305.
- [3] Cotton CM. *Ethnobotany: Principle and application*. New York: John Wiley and Sons; 1996, p. 399.
- [4] Poonam K, Singh GS. Ethnobotanical study of medicinal plants used by the Taungya community in Terai Arc Landscape. *Indian J Ethnopharmacol* 2009; **123**:167–176.
- [5] Rao RR. Ethnobiology in India: some key issues on traditional knowledge and sustainable development. In: *Ethnobiology and conservation of cultural and biological diversity: proceedings of the Fifth International Congress of Ethnobiology*, Nairobi, Kenya, September 2–6 1996. Kenya: National Museums of Kenya; 1996, p. 2.
- [6] Kala CP, Sajwan BS. Revitalizing Indian systems of herbal medicine by the National Medicinal Plants Board through institutional networking and capacity building. *Curr Sci* 2007; **93**: 797–806.
- [7] Ahmad SS, Husain SZ. Ethno medicinal survey of plants from salt range (Kallar Kahar) of Pakistan. *Pak J Bot* 2008; **40**: 1005–1011.
- [8] Tovey PA, Broom AF, Chatwin J, Ahmad S, Hafeez M. Use of traditional, complementary and allopathic medicines in Pakistan by cancer patients. *Rural Remote Health* 2005; **5**: 447.
- [9] Khan MA, Qaiser M. Halophytes of Pakistan: characteristics, distribution and potential economic usages. In: Khan MA, editor. *Sabkha ecosystems Volume II: West and Central Asia*. Netherlands: Springer; 2006, p. 129–153.
- [10] Qasim M, Gulzar S, Shinwari ZK, Aziz I, Khan MA. Traditional ethnobotanical uses of halophytes from Hub, Balochistan. *Pak J Bot* 2010; **42**: 1543–1551.
- [11] Qasim M, Gulzar S, Khan MA. Halophytes as medicinal plants. In: Ozturk M, Mermut AR, Celik A, editors. *Urbanisation, land use, land degradation and environment*. Delhi: Daya Publishing House; 2011, p. 330–343.
- [12] Abideen Z, Ansari R, Khan MA. Halophytes: Potential source of ligno-cellulosic biomass for ethanol production. *Biomass Bioenerg* 2011; **35**: 1818–1822.
- [13] Khan MA, Ansari R, Ali H, Gul B, Nielsen BL. *Panicum turgidum*, a potentially sustainable cattle feed alternative to maize for saline areas. *Agri Ecosys Environ* 2009; **129**: 542–546.
- [14] Masters DG, Benes SE, Norman HC. Biosaline agriculture for

- forage and livestock production. *Agri Ecosys Environ* 2007; **119**: 234–248.
- [15] Rogers ME, Craig AD, Munns R, Colmer TD, Nichols PGH, Malcolm CV, et al. The potential for developing fodder plants for the salt-affected areas of southern and eastern Australia: an overview. *Aus J Exp Agric* 2005; **45**, 301–329.
- [16] Midgley SJ, Turnbull JW. Domestication and use of Australian Acacias: case studies of five important species. *Aust Syst Bot* 2003; **16**: 89–102.
- [17] Martin GJ. *Ethnobotany: A methods manual*. London, UK: Chapman and Hall; 1995.
- [18] Khan SM, Page S, Ahmad H, Shaheen H, Ullah Z, Ahmad M, et al. Medicinal flora and ethnoecological knowledge in the Naran Valley, Western Himalaya, Pakistan. *J Ethnobiol Ethnomed* 2013; **9**: 4.
- [19] Stepp JR. The role of weeds as sources of pharmaceuticals. *J Ethnopharmacol* 2004; **92**: 163–166.
- [20] Albuquerque UP, Hanazaki N. Five problems in current ethnobotanical research and some suggestions for strengthening them. *Human Ecol* 2009; **37**:653–661.
- [21] Voeks RA. Disturbance pharmacopoeias: medicine and myth from the humid tropics. *Ann Assoc Am Geogr* 2004; **94**: 868–888.
- [22] Abdul MM, Sarker AA, Saiful IM, Muniruddin A. Cytotoxic and antimicrobial activity of the crude extracts of *Abutilon indicum*. *Int J Pharmaco Phytochem Res* 2010; **2**: 1–4.
- [23] Saini ML, Saini R, Roy S, Kumar A. Comparative pharmacognostical and antimicrobial studies of *Acacia* species (Mimosaceae). *J Med Plant Res* 2008; **2**: 378–386.
- [24] Londonkar R, Reddy CV, Kumar K. Potential antibacterial and antifungal activity of *Achyranthes aspera* L. *Recent Res Sci Technol* 2011; **3**: 53–57.
- [25] Kumar D, Prasad DN, Bhatnagar SP. Comparison of diuretic activity of ethanolic extract of *Aerva lanata* (Linn.) Juss. Ex. Schult & *Aerva tomentosa* Forsk. Family: Amaranthaceae. *Anc Sci Life* 2005; **25**: 66–68.
- [26] Sethi A, Sharma RA. Antioxidant activity with total phenolic constituents from *Aerva tomentosa* forsk. *Int J Pharma Biosci* 2011; **2**: 596–603.
- [27] Atta AH, Kenawy S, Awaad A, Melegy RE. Anti-Inflammatory, antipyretic and antioxidant effect of some medicinal plant extracts. *J Comp Integr Med* 2011; **8**: doi: 10.2202/1553–3840.1086.
- [28] Bencheraiet B, Kherrab H, Kabouche A, Kabouche Z, Jay M. Flavonols and antioxidant activity of *Ammi visnaga* L. (Apiaceae). *Record Nat Prod* 2011; **5**: 52–55.
- [29] Yurina EV, Yurina TP, Lekomtseva SN, Karavaev VA, Solntsev MK. Antifungal activity of aqueous extracts from *Artemisia scoparia* and *Rumex confertus*. Modern fungicides and antifungal compounds IV: 14th International Reinhardsbrunn Symposium, Friedrichroda, Thuringia, Germany, April 25–29, 2004. 2004, p. 277–280.
- [30] Negahban M, Moharramipour S, Sefidkon F. Chemical composition and insecticidal activity of *Artemisia scoperte* essential oil against three coleopteran stored-product insects. *J Asia-Pac Entomol* 2006; **9**: 381–388.
- [31] Bandaranayake WM. Traditional and medicinal uses of mangroves. *Mangroves Salt Marshes* 1998; **2**: 133–148.
- [32] Nahak G, Sahu RK. *In vitro* antioxidative activity of *Azadirachta indica* and *Melia azedarach* leaves by DPPH scavenging assay. *J Am Sci* 2010; **6**: 123–128.
- [33] Simin K, Khaliq-Uz-Zaman SM, Ahmad VU. Antimicrobial activity of seed extracts and bordenolide from *Caesalpinia bonduc* (L.) Roxb. *Phytother Res* 2001; **15**: 437–440.
- [34] Yesmin MN, Uddin SN, Mubassara S, Akond MA. Antioxidant and antibacterial activities of *Calotropis procera* Linn. *Am Eurasian J Agric Environ Sci* 2008; **4**: 550–553.
- [35] Pirezada AJ, Shaikh W, Maka GA, Shah SIS, Mughal S. Anti-fungal activity of different solvent extracts of medicinal plants *Capparis decidua* Edgew and *Salvadora persica* Linn. against different parasitic fungi. *Pak J Agri Agri Eng Vet Sci* 2009; **25**: 26–34.
- [36] Sharma B, Kumar P. Extraction and pharmacological evaluation of some extracts of *Tridax procumbens* and *Capparis decidua*. *Int J Appl Res Nat Prod* 2008; **1**: 5–12.
- [37] Ahmed S, Qureshi S, Kapadia Z, Badar Y. Cathartic activity of *Cassia* species. *Pak J Pharm Sci* 1989; **2**: 37–45.
- [38] Nayak DP, Swain PK, Panda OP, Pattanaik P, Srinivas B. Antimicrobial and anthelmintic evaluation of *Chenopodium album*. *Int J Pharma World Res* 2010; **1**: 1–15.
- [39] Dallak M, Bin-Jalial. Antioxidant activity of *Citrullus colocynthis* pulp extract in the Rbc's of alloxan-induced diabetic rats. *Pak J Physiol* 2010; **6**: 1–5.
- [40] Marzouk BA, Marzouk BZ, Fenina BN, Bouraoui CA, Aoumi AM. Anti-inflammatory and analgesic activities of Tunisian *Citrullus colocynthis* Schrad. immature fruit and seed organic extracts. *Eur Rev Med Pharmacol Sci* 2011; **15**: 665–672.
- [41] Chandak RR, Bhairat NK, Devdhe SJ, Majmudar HF. *In vitro* evaluation of anthelmintic potential of leaves of *Cleome viscosa* Linn. *Int J Pharm Sci Rev Res* 2010; **5**: 77–79.
- [42] Vohora SB, Shamsi MA, Khan MSY. Antipyretic and analgesic studies on the diacetate of a new triterpenic acid isolated from *Corchorus depressus* L. *J Ethnopharmacol* 1981; **4**: 223–228.
- [43] Sunita P, Jha S, Pattanayak SP. Anti-inflammatory and *in-vivo* antioxidant activities of *Cressa cretica* Linn., a halophytic plant. *Middle-East J Sci Res* 2011; **8**: 129–140.
- [44] Alam K, Hussain T, Ahmad S. *Antipyretic activity of medicinal plants of Cholistan Desert Pakistan: Evaluation of antipyretic activity of some medicinal plants from Cholistan Desert Pakistan*. Saarbrücken, Germany: Lambert Academic Publishing; 2011.
- [45] Nagulendran K, Velavan S, Mahesh R, Begum VH. *In vitro* antioxidant activity and total polyphenolic content of *Cyperus rotundus* rhizomes. *Electron J Chem* 2007; **4**: 440–449.
- [46] Kumar KA, Sharvanee PS, Choudhary RK. Chemical composition and antimicrobial activity of the essential oil of *Desmostachya bipinnata* Linn. *Int J Phytomed* 2010; **2**: 436–439.
- [47] Gami B, Parabia F. Screening of methanol and acetone extract for antimicrobial activity of some medicinal plants species of Indian folklore. *Int J Res Pharm Sci* 2011; **2**: 69–75.
- [48] Patel MB, Mishra SH. Hypoglycemic activity of C-glycosyl flavonoid from *Enicostemma hyssopifolium*. *Pharm Biol* 2011; **49**:

- 383–391.
- [49] Babayi H, Kolo I, Okogun JI, Ijah UJJ. The antimicrobial activities of methanolic extracts of *Eucalyptus camaldulensis* and *Terminalia catappa* against some pathogenic microorganisms. *Biokemistri* 2004; **16**:106–111.
- [50] Medhi SM, Reza SA, Mahnaz K, Reza AM, Hadjiakhoondi A, Fatemeh M, et al. Phytochemistry and larvicidal activity of *Eucalyptus camaldulensis* against malaria vector, *Anopheles stephensi*. *Asia Pac J Trop Med* 2010; **3**: 841–845.
- [51] Dahot MU, Khan MY, Memon AN. Screening of some Pakistani plants for milk clotting activity. *J Islamic Acad Sci* 1990; **3**: 284–286.
- [52] Ogueke CC, Ogbulie JN, Okoli IC, Anyanwu BN. Antibacterial activities and toxicological potentials of crude ethanolic extracts of *Euphorbia hirta*. *J Am Sci* 2007; **3**: 11–16.
- [53] Soomro AL, Jafarey NA. Effect of *Fagonia indica* on experimentally produced tumours in rats. *J Pak Med Assoc* 2003; **53**(6): 440–441.
- [54] Sharma S, Joseph L, George M, Gupta V. Analgesic and antimicrobial activities of *Fagonia indica*. *Pharmacologyonline* 2009; **3**: 623–632.
- [55] Saadabi AMA, Moglad EH. Experimental evaluation of certain sudanese plants used in folkloric medicine for their antibacterial activity (*In-vitro* tests). *J Appl Sci Res* 2011; **7**: 253–256.
- [56] Dahot MU. Antibacterial and antifungal activity of small protein of *Indigofera oblongifolia* leaves. *J Ethnopharmacol* 1999; **64**: 277–282.
- [57] Upwar N, Patel R, Waseem N, Mahobia NK. Anti-inflammatory potential of alcoholic extract of *Indigofera oblongifolia* Forsk. *Int J Res Pharm Sci* 2011; **2**: 23–25.
- [58] Dipak AT, Tejal BC, Sanjay RC. Wound healing activity of *Indigofera cordifolia* aerial part extracts in rats. *J Pharm Res* 2010; **3**: 417–419.
- [59] Ashraf A, El-Bassuony NM, Hamid A. Antibacterial coumarins isolated from *Launaea resedifolia*. *Chem Plant Raw Mater* 2006; **1**: 65–68.
- [60] Fatimi MA, Wurster M, Schroder G, Lindequist U. Antioxidant, antimicrobial and cytotoxic activities of selected medicinal plants from Yemen. *J Ethnopharmacol* 2007; **111**: 657–666.
- [61] Shah KA, Mandev BP, Shreya SS, Kajal NC, Parul KP, Natavarlal MP. Antihyperlipidemic activity of *Mangifera indica* L. leaf extract on rats fed with high cholesterol diet. *Der Pharmacia Sinica* 2010; **1**: 156–161.
- [62] Leite ACR, Araujo TG, Carvalho BD, Maia MBS, Lima VLD. Characterization of the antidiabetic role of *Parkinsonia aculeata* (Caesalpinaceae). *Evid Based Complement Alternat Med* 2011; **2011**(2011): 1–9.
- [63] Ahmida MH. Evaluation of *in vivo* antioxidant and hepatoprotective activity of *Portulaca oleracea* L. against paracetamol-induced liver toxicity in male rats. *Am J Pharm Toxicol* 2010; **5**: 167–176.
- [64] Sharma N, Garg V, Paul A. Antihyperglycemic, antihyperlipidemic and antioxidative potential of *Prosopis cineraria* bark. *Indian J Clin Biochem* 2010; **25**: 193–200.
- [65] Ahmad A, Khan KA, Ahmad V. Immunomodulating effect of Juliflorine on the antibody response to *Listeria hemolysin* preliminary report. *J Islamic Acad Sci* 1992; **5**: 189–193.
- [66] Bentes J, Amaral-da VD, Rita A, Santos CD, Mendes-da AM, Silva ED, et al. Cytotoxicity effect of alkaloidal extract from *Prosopis juliflora* Sw. D.C. (Algaroba) pods on glial cells. *Braz J Vet Res Anim Sci* 2006; **43**: 50–58.
- [67] Orak HH, Demirci AS, Orak TG. Antibacterial and antifungal activity of pomegranate (*Punica granatum* L.) peel. *Electron J Environ Agric Food Chem* 2011; **10**: 1958–1969.
- [68] Kotamballi N, Murthy C. Studies on antioxidant activity of pomegranate (*Punica granatum*) peel extract using *in vivo* models. *J Agric Food Chem* 2002; **50**: 4791–4795.
- [69] Yadav JP, Saini S, Kalia AN, Dangi AS. Hypoglycemic and hypolipidemic activity of ethanolic extract of *Salvadora oleoides* in normal and alloxan-induced diabetic rats. *Indian J Pharmacol* 2008; **40**: 23–27.
- [70] Nawal AK, Al-Sabawi, Khalik AK, Abdal AS, Taha MY. The antimicrobial activity of *Salvadora persica* solution (miswak-siwak) as root canal irrigant (a comparative study). *J Pure Appl Sci* 2007; **4**: 69–91.
- [71] Sheeba E. Antibacterial activity of *Solanum surattense* burm. f. *Kathmandu Univ J Sci Eng Technol* 2010; **6**: 1–4.
- [72] Kumar A, Ilavarasan R, Jayachandran T, Deecaraman M, Aravindan P, Padmanabhan N, et al. Anti-diabetic activity of *Syzygium cumini* and its isolated compound against streptozotocin-induced diabetic rats. *J Med Plant Res* 2008; **2**: 246–249.
- [73] Bobbarala V, Vadlapudi VR, Naidu KC. *In vitro* antimicrobial properties of mangrove plant *Thespesia populneoides* against selected clinical, plant and aquatic pathogens. *Biotechnol Res Asia* 2009; **6**: 351–354.
- [74] Sunder AS, Reddy ARN, Kiran G, Thirumurugu S. Antihyperlipidemic and antioxidant activity of methanolic extract of *Trianthema portulacastrum* in rats fed a high-fat diet. *J Herbs Spices Med Plant* 2010; **16**: 193–202.
- [75] Al-Bayati FA, Al-Mola HF. Antibacterial and antifungal activities of different parts of *Tribulus terrestris* L. growing in Iraq. *J Zhejiang Univ Sci B* 2008; **9**: 154–159.
- [76] Singh S, Gupta YK. Aphrodisiac activity of *Tribulus terrestris* Linn. in experimental models in rats. *J Men Health* 2011; **8**: S75–S77.
- [77] Singh PS, Vidyasagar GM. Ethnomedicinal plants used in the treatment of skin diseases in Hyderabad Karnataka region, Karnataka, India. *Asia Pac J Trop Biomed* 2013; **3**: 882–886.
- [78] Panchawat S. *In vitro* free radical scavenging activity of leaves extracts of *Withania somnifera*. *Recent Res Sci Technol* 2011; **3**: 40–43.
- [79] Jeyaprakash K, Ayyanar M, Geetha KN, Sekar T. Traditional uses of medicinal plants among the tribal people in Theni District (Western Ghats), Southern India. *Asia Pac J Trop Biomed* 2011; **1**: S20–S25.
- [80] Chanda S, Kaneria M, Vaghasiya YK. Evaluation of antimicrobial potential of some Indian medicinal plants against some pathogenic microbes. *Indian J Nat Prod Resour* 2011; **2**: 225–228.