

Journal of Coastal Life Medicine

journal homepage: www.jclmm.com



Document heading doi: 10.12980/JCLM.2.2014APJTD-2014-0095 © 2014 by the Journal of Coastal Life Medicine. All rights reserved.

Prevalence of *Helicobacter pylori cagA* genotype among dyspeptic patients in Southern ThailandSueptrakool Wisessombat^{1*}, Chatruthai Meethai²¹School of Allied Health Sciences and Public Health, Walailak University, Tha Sala, Nakhon Si Thammarat Province, Thailand²Faculty of Medical Technology, Prince of Songkla University, Hat Yai, Songkhla Province, Thailand

PEER REVIEW

Peer reviewer

Assoc. Prof. Dr. Siroj Jitsuronk, Faculty of Medical Technology, Prince of Songkla University, Hat Yai, Songkhla, Thailand.

E-mail: siroj.jit@gmail.com

Comments

This article is interesting and revealed that in Southern Thailand, the prevalence of *cagA* gene in *H. pylori* was found significantly high in female. Details on Page 897

ABSTRACT

Objective: To investigate the prevalence of *Helicobacter pylori* (*H. pylori*) infection in dyspepsia patients and its relation to virulence factor *cagA* gene.

Methods: In total, 110 gastric biopsies from dyspeptic patients were comparatively studied using rapid urease test and multiplex polymerase chain reaction (PCR).

Results: Multiplex PCR detected three genes of 16S rRNA, *cagA*, and *ureC*. *H. pylori* was detected in 14 gastric biopsies (13%). Significantly higher numbers of female were infected. Furthermore, *cagA* gene was found in all *H. pylori*-positive specimens. In addition, the result indicated that the multiplex PCR with annealing temperature at 57 °C was able to effectively amplify specific products.

Conclusions: The results confirmed high prevalence of *cagA* gene in *H. pylori* among dyspeptic patients in Southern Thailand.

KEYWORDS

Helicobacter pylori, Multiplex polymerase chain reaction, Dyspepsia, *cagA* gene

1. Introduction

Helicobacter pylori (*H. pylori*), a Gram-negative microaerobic bacterium, is associated with human gastritis, gastric ulcer and gastric cancer^[1]. Cytotoxin associated gene *cagA* is one of the most studied virulence factors of *H. pylori*. *cagA* has been proposed as a marker for a genomic pathogenicity island^[2]. *H. pylori cagA*-positive

strains have been observed to be more virulent than the *H. pylori cagA*-negative strains. The *cagA*-positive strain increases the risk of development of atrophic gastritis, mucosal inflammation, and adenocarcinoma^[3].

Histology has been considered to be the gold standard for detection of *H. pylori*. However, the detection of *H. pylori* relies upon a number of gastric biopsies, staining methods, and the level of experience of the examining

*Corresponding author: Sueptrakool Wisessombat, School of Allied Health Sciences and Public Health, Walailak University, Tha Sala, Nakhon Si Thammarat Province, Thailand.

Tel: +66 75672472

Fax: +66 75672106

E-mail: sueptrakool.wi@wu.ac.th

Foundation Project: Supported by Prince of Songkla University, Research Development Fund, Fiscal Year 2011–2012.

Article history:

Received 28 Mar 2014

Received in revised form 6 Apr, 2nd revised form 13 Apr, 3rd revised form 18 Apr 2014

Accepted 20 May 2014

Available online undetermined

pathologist^[4]. Molecular methods based on polymerase chain reaction (PCR) amplification are rapid, specific and sensitive. A number of PCR-based methods have been reported for the detection of *Helicobacter*^[5–7]. In Southern Thailand, the epidemiological studies on prevalence of *H. pylori* infection are very few.

The objective of the present study was to investigate the prevalence of *H. pylori* infection among dyspeptic patients in Southern Thailand. We also established a multiplex PCR for the identification of *H. pylori*. In addition, *cagA* gene-based multiplex PCR can simultaneously detect the presence of *cagA* gene which is responsible for pathogenesis of *H. pylori* infection.

2. Materials and methods

2.1. Bacterial strains and culture conditions

Reference strains including *H. pylori* NCTC 11637 and *H. pylori* NCTC 11638 were used for development of a multiplex PCR. *Helicobacter* species were cultured on Brucella blood agar (BBL, USA) with 10% defibrinated horse blood (Oxoid, UK). Plates were incubated at 37 °C for 48 h under microaerobic atmosphere using gas pack system (Oxoid).

2.2. Gastric biopsies

Gastric biopsies were collected from Institute of Gastroenterology and Hepatology, Songklanagarind Hospital, Prince of Songkla University, Thailand. A total of 110 dyspeptic patients undergoing upper endoscopy were biopsied and tested for *H. pylori* infection by a *Campylobacter*-like organism (CLO) test (Kimberly-Clark, USA) and multiplex PCR. The CLO test was performed according to the manufacturer's instructions, and the results were interpreted after 24 h.

2.3. Multiplex PCR

Bacterial DNA was extracted and purified directly from biopsy specimens by QIAamp DNA Mini Kit (QIAamp, USA). The identification of *H. pylori* confirmed specific primers. In this study, a multiplex PCR was designed to detect three genes of 16S rRNA, *cagA* encoding for virulence factor cytotoxin associated gene A, and *ureC* for housekeeping urease gene C (Table 1). PCR was performed in a total reaction volume of 25 mL containing 1× TopTaq Master (QIAamp), 1.5 mmol/L MgCl₂, 200 mmol/L deoxynucleotide

triphosphates, 1.25 IU *Taq* polymerase, 20 μmol 16S rRNA primers for *H. pylori*, 15 μmol each of *cagA* primers and *ureC* primers for *H. pylori*. Amplification consisted of initial denaturation at 94 °C for 4 min, followed by denaturation at 94 °C for 30 seconds, primers annealing at 50–60 °C for 30 seconds, and extension at 72 °C for 30 seconds. The samples were amplified for 40 cycles, with a final extension step at 72 °C for 5 min. PCR cycles were carried out in PTC-100, Peltier Thermal Cycler (Pegasus Scientific, USA). About 2 μL amplified products were analysed by 2% agarose (Gibco-BRL Life Technologies, USA) gel electrophoresis in Tris-Acetate-EDTA buffer at 100 V for 35 min. PCR products were visualized after ethidium bromide staining.

Table 1

Primers used in this study.

Target gene	Primers sequences	Annealing temperature (°C)	Amplicon size (base pair)	References
16S rRNA	F 5' TAA GAG ATC AGC CTA TAT GTC C 3' R 5' TCC CAC GCT TTA AGC GCA AT 3'	56	534	[22]
<i>cagA</i>	F 5' AAT ACA CCA ACG CCT CCA AG 3' R 5' TTG TTG CCG CTT TTG CTC TC 3'	59	400	[23]
<i>ureC</i>	F 5' AAG CTT TTA GGG GTG TTA GGG GTT 3' R 5' AAG CTT ACT TTC TAA CAC TAA CGC 3'	57	294	[24]

2.4. Statistical analysis

Data were subjected to analysis of invariance. Determination of the prevalence of *H. pylori* infection in relation to gender and age were carried out by Fisher's exact test (2-tailed test). Statistical analysis was performed using the Statistical Package for Social Sciences package version 12.0 (SPSS, USA).

3. Results

In total, in 110 dyspeptic patients 56 were female and 54 were male. *H. pylori* infected patients were evaluated for the relation of gender and age as shown in Table 2.

Table 2

H. pylori infection rates in relation to gender and age.

Gender	Age (years old)	Total	<i>H. pylori</i> positive (%)
Female	<20	2	0
	21–40	3	0
	41–60	25	3 (12%)
	>60	26	7 (27%)
	Total	56	10 (18%)
Male	<20	2	0
	21–40	2	0
	41–60	22	2 (9%)
	>60	28	2 (7%)
	Total	54	4 (7%)

The results demonstrated that *H. pylori* infection rates were significantly higher ($P < 0.05$) in female aged over 60 years.

The presence of *H. pylori* in the gastric biopsies was detected by CLO test and PCR. The results showed that *H. pylori* were positive in 14 gastric biopsies (13%). *cagA* gene was detected in all *H. pylori*-infected dyspeptic patients. Moreover, the optimal condition of the multiplex PCR was carried out with a single tube method by incorporating all specific primers. The combination of 16S rRNA, *cagA*, and *ureC* primers were able to be detected at 57 °C annealing temperatures (data not shown).

4. Discussion

It has been showed that *H. pylori* infection rate in dyspeptic patients was 13%. Nevertheless, the prevalence of *H. pylori cagA* genotype was 100%. Likewise, the positive rate for the *cagA* gene in *H. pylori* of dyspeptic patients was 94% in Northeast Thailand[8]. Whereas, it was reported that the prevalence of *cagA* gene was found to be 60%–70% in Western countries[9].

In Thailand, *H. pylori* infection rate was 34.1%[10]. Moreover, 48% of dyspeptic patients were infected with *H. pylori*[11–13]. Similarly, the prevalence of *H. pylori* infection changes considerably with age[14,15].

The *H. pylori cagA* genotype strains are associated with gastric carcinogenesis by increasing interleukin 8 secretion, NF- κ B activation, and stimulation of cell proliferation[2,16,17]. The prevalence of gastric cancer in Thailand was reported to be lower than that in other South–East Asia countries even the prevalence of *H. pylori* infection was higher[18]. In Thailand, the prevalence of gastric cancer was 1.5%, while, it was 3.3% in Malaysia[19]. Furthermore, the Western type *cagA* was detected to be more frequently than the East Asian type in Thai dyspeptic patients. It was also found significantly more common in patients with a gastric ulcer but was not significant in gastric cancer[20]. Recent study have revealed that the variation of Western type *cagA* gene may be involved in the development of diseases[21].

In conclusion, these observations indicated that the *cagA* gene is an important virulence factor for *H. pylori*-infected dyspepsia patients. In addition, our multiplex PCR has allowed simultaneous amplification of *H. pylori* virulent genes direct from biopsies.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgements

This work was supported by Prince of Songkla University, Research Development Fund, Fiscal Year 2011–2012. We also thank Faculty of Medical Technology, Prince of Songkla University for the use of their lab facilities.

Comments

Background

H. pylori is a Gram-negative bacterium causing human gastritis, gastric ulcer and gastric cancer. Cytotoxin associated gene A is one of the most studied virulence factors of *H. pylori*. *H. pylori cagA*-positive strains have been observed to be more virulent than the *H. pylori cagA*-negative strains. Diagnosis of *H. pylori cagA* genotype infections can be performed by PCR.

Research frontiers

This study was performed in order to determine the prevalence of *H. pylori* infection among dyspeptic patients in Southern Thailand. Furthermore, also this study established a new multiplex PCR for the identification of *H. pylori* using *cagA* gene-based PCR.

Related reports

The manuscript discussed the prevalence of *H. pylori* infections from the United States and Southeast Asia. To establish a new multiplex PCR, the gastric biopsies from dyspeptic patients were comparatively studied using CLO test and multiplex PCR.

Innovations and breakthroughs

This study indicated that the multiplex PCR with annealing temperature at 57 °C was able to effectively amplify specific PCR products. Moreover, the *cagA* gene-based PCR has allowed simultaneous directly detection of virulent genes from gastric biopsies.

Applications

It may be significant to know the distribution of *cagA*

gene in dyspeptic patients. The results of the present study confirmed the high prevalence of *cagA* gene in *H. pylori* among dyspeptic patients in Southern Thailand, especially in female aged over 60 years.

Peer review

This article is interesting and revealed that in Southern Thailand, the prevalence of *cagA* gene in *H. pylori* was found significantly high in female.

References

- [1] Kusters JG, van Vliet AH, Kuipers EJ. Pathogenesis of *Helicobacter pylori* infection. *Clin Microbiol Rev* 2006; **19**(3): 449–490.
- [2] Noto JM, Peek RM Jr. The *Helicobacter pylori* *cag* pathogenicity island. *Methods Mol Biol* 2012; **921**: 41–50.
- [3] Kaklikkaya N, Cubukcu K, Aydin F, Bakir T, Erkul S, Tosun I, et al. Significance of *cagA* status and *vacA* subtype of *Helicobacter pylori* in determining gastric histopathology: virulence markers of *H. pylori* and histopathology. *J Gastroenterol Hepatol* 2006; **21**: 1042–1047.
- [4] Cirak MY, Akyon Y, Megraud F. Diagnosis of *Helicobacter pylori*. *Helicobacter* 2007; **12**: 4–9.
- [5] Khademi F, Faghri J, Moghim S, Esfahani BN, Fazeli H, Poursina F, et al. The study of mutation in 23S rRNA resistance gene of *Helicobacter pylori* to clarithromycin in patients with gastrointestinal disorders in Isfahan–Iran. *Adv Biomed Res* 2014; **3**: 98.
- [6] Mesquita B, Gonçalves MJ, Pacheco P, Lopes J, Salazar F, Relvas M, et al. *Helicobacter pylori* identification: a diagnostic/confirmatory method for evaluation. *Curr Microbiol* Forthcoming 2014.
- [7] Morimoto N, Takeuchi H, Nishida Y, Morisawa M, Yoshikawa T, Morita T, et al. Clinical application of the diversilab microbial typing system using repetitive sequence–based PCR for characterization of *Helicobacter pylori* in Japan. *J Clin Lab Anal* 2014; doi: 10.1002/jcla.21758.
- [8] Chomvarin C, Phusri K, Sawadpanich K, Mairiang P, Namwat W, Wongkham C, et al. Prevalence of *cagA* EPIYA motifs in *Helicobacter pylori* among dyspeptic patients in northeast Thailand. *Southeast Asian J Trop Med Public Health* 2012; **43**: 105–115.
- [9] Alm RA, Ling LS, Moir DT, King BL, Brown ED, Doig PC, et al. Genomic–sequence comparison of two unrelated isolates of the human gastric pathogen *Helicobacter pylori*. *Nature* 1999; **397**: 176–180.
- [10] Vilaichone RK, Gumnarai P, Ratanachu–Ek T, Mahachai V. Nationwide survey of *Helicobacter pylori* antibiotic resistance in Thailand. *Diagn Microbiol Infect Dis* 2013; **77**: 346–349.
- [11] Atisook K, Kachinthorn U, Luengrojanakul P, Tanwadee T, Pakdirat P, Puapairoj A. Histology of gastritis and *Helicobacter pylori* infection in Thailand: a nationwide study of 3776 cases. *Helicobacter* 2003; **8**: 132–141.
- [12] Chomvarin C, Kulsuntiwong P, Mairiang P, Sangchan A, Kulabkhow C, Chau–in S, et al. Detection of *Helicobacter pylori* in dyspeptic patients and correlation with clinical outcomes. *Southeast Asian J Trop Med Public Health* 2005; **36**: 917–922.
- [13] Mitipat N, Siripermpool P, Jadwattanakul T, Chaunthongkum S. The prevalence of *Helicobacter pylori* infection in patients with gastrointestinal symptoms in Chon Buri, Thailand. *Southeast Asian J Trop Med Public Health* 2005; **36**: 341–346.
- [14] Sasidharan S, Lachumy SJ, Ravichandran M, Latha LY, Gegu SR. Epidemiology of *Helicobacter pylori* among multiracial community in Northern Peninsular, Malaysia: effect of age across race and gender. *Asian Pac J Trop Med* 2011; **4**: 72–75.
- [15] Hunt RH, Xiao SD, Megraud F, Leon–Barua R, Bazzoli F, van der Merwe S, et al. *Helicobacter pylori* in developing countries. World gastroenterology organisation global guideline. *J Gastrointest Liver Dis* 2011; **20**: 299–304.
- [16] Boonyanugomol W, Chomvarin C, Hahnvajanawong C, Sripa B, Kaparakis–Liaskos M, Ferrero RL. *Helicobacter pylori* *cag* pathogenicity island (*cagPAI*) involved in bacterial internalization and IL–8 induced responses via NOD1–and MyD88–dependent mechanisms in human biliary epithelial cells. *PLoS One* 2013; doi: 10.1371/journal.pone.0077358.
- [17] Parsonnet J, Friedman GD, Orentreich N, Vogelstein H. Risk for gastric cancer in people with *cagA* positive or *cagA* negative *Helicobacter pylori* infection. *Gut* 1997; **40**: 297–301.
- [18] Singh K, Ghoshal UC. Causal role of *Helicobacter pylori* infection in gastric cancer: an Asian enigma. *World J Gastroenterol* 2006; **12**: 1346–1351.
- [19] Rahman R, Asombang AW, Ibdah JA. Characteristics of gastric cancer in Asia. *World J Gastroenterol* 2014; **20**: 4483–4490.
- [20] Chomvarin C, Namwat W, Chaicumpar K, Mairiang P, Sangchan A, Sripa B, et al. Prevalence of *Helicobacter pylori* *vacA*, *cagA*, *cagE*, *iceA* and *babA2* genotypes in Thai dyspeptic patients. *Int J Infect Dis* 2008; **12**: 30–36.
- [21] Boonyanugomol W, Chomvarin C, Sripa B, Chau–in S, Pughkhem A, Namwat W, et al. Molecular analysis of *Helicobacter pylori* virulent–associated genes in hepatobiliary patients. *HPB (Oxford)* 2012; **14**: 754–763.
- [22] Tiwari SK, Khan AA, Ibrahim M, Habeeb MA, Habibullah CM. *Helicobacter pylori* and other *Helicobacter* species DNA in human bile samples from patients with various hepato–biliary disease. *World J Gastroenterol* 2006; **12**: 2181–2186.
- [23] Russo F, Notarnicola M, Di Matteo G, Leoci C, Caruso ML, Pirrelli M, et al. Detection of *Helicobacter pylori* *cagA* gene by polymerase chain reaction in faecal samples. *Eur J Gastroenterol Hepatol* 1999; **11**: 251–256.
- [24] Bickley J, Owen RJ, Fraser AG, Pounder RE. Evaluation of the polymerase chain reaction for detecting the urease C gene of *Helicobacter pylori* in gastric biopsy samples and dental plaque. *J Med Microbiol* 1993; **39**: 338–344.