

# Journal of Coastal Life Medicine

journal homepage: [www.jclmm.com](http://www.jclmm.com)



Document heading doi: 10.12980/JCLM.3.201514J97

©2015 by the Journal of Coastal Life Medicine. All rights reserved.

## Stress reaction in freshwater fish in response to extreme impacts and during the reproduction period

Vladimir Ivamovich Martemyanov\*

*Papanin Institute for the Biology of Inland Waters, Russian Academy of Sciences, 152742 Borok, Yaroslavl oblast, Russia*

### ARTICLE INFO

#### Article history:

Received 20 Dec 2014

Received in revised form 26 Dec 2014

Accepted 28 Jan 2015

Available online 5 Feb 2014

#### Keywords:

Fish

Stress

Spawning

Reproduction

Stages of alarm

Resistance

Exhaustion

Protective functions

Negative processes

### ABSTRACT

The original and published data on the physiological and biochemical reactions in fish in response to stress impact and during the reproduction period are presented. It is shown that at the initial period of stress and before spawning the protective functions in fish are enhanced which relates to the adaptation syndrome. However, during the initial period of stress the damaging effects prevail over the adaptive processes resulting in the decrease in the level of resistance ("alarm reaction" according to Selye). In contrary, during pre-spawning period protective systems dominate along with enhancement of general adaptation syndromes. This facilitates the increase in the level of resistance in the parental fish ("resistance stage" according to Selye). Before the spawning the alarm stage is not manifested. The state comparable to strong stress occurs in the parental fish only during the spawning. This state is accompanied by decrease in the resistance of organism in spawner evidencing exhaustion of protective functions ("exhaustion stage" according to Selye). With time, both at stress and after spawning, within two to three weeks period, the physiological-biochemical parameters recover and stabilize with the normal limits.

## 1. Introduction

Several researchers have noted that during the reproduction period parental fish exhibit the state resembling stress[1-4]. It is unclear if the stress in reproducing fish identical to the stress in response to extreme factors or are there any differences between these states? Clarification of this uncertainty necessitate the comparative analysis of physiological and biochemical processes observed in fish in response to stressors and during the period of reproduction.

\*Corresponding author: Vladimir Ivamovich Martemyanov, Papanin Institute for the Biology of Inland Waters, Russian Academy of Sciences, 152742 Borok, Yaroslavl oblast, Russia.

E-mail: [martem@ibiw.yaroslavl.ru](mailto:martem@ibiw.yaroslavl.ru)


Foundation Project: Supported by the budgetary state project: Studying of adaptations of organisms, populations, communities of aquatic organisms and modeling ecosystems to environmental factors. State registration No. 01201351909.

## 2. Dynamics of physiological and biochemical parameters in freshwater fish from the beginning of stress impact to the end of acclimation to new conditions

The comparative analysis of available data revealed the consequence of the processes taking place in freshwater fish in response to stress impacts[5]. It was shown that on the one hand the stressors cause a series of damages in the organism (Figure 1, right part), but on the other hand, enhance the protective functions (Figure 1, left part) withstanding development of harmful processes.

### 2.1. Damaging effects of stress-factors

Many factors of physical and chemical (toxicants) natures cause direct structural damages, first of fish gills[6]. However, all types of




**Figure 1.** Schematic diagram of the conjugated processes related to the GAS (left part) and negative symptoms (right part) during the stress[5]. MSH: melanophor-stimulating hormone.

stressors possess common mechanism of destabilization of organism. In 1972 it was hypothesized that damaging and malfunctions at stress are realized via peroxides and free radicals[7]. The stressors reinforce nonenzymatic pathway of the oxidation by free radicals of energetic and plastic substances in the organs and tissues[8]. This results in the formation of aggressive free radicals. These radicals directly bind with oxygen and produce toxic peroxidation products.

It was shown that during initial stage of stress the organs and tissues of mammals[7,9,10], and fish[11-15] accumulate harmful substances possessing hemolytic activity: aggressive radical, "debris of organic molecules, lipid peroxides, etc. These substances cause various damaged in the organism (Figure 1, right part), including

increase in the permeability of cellular membranes. As a result, the blood passing fish gill capillary system loses ions, mainly sodium and chlorine[16-21] moving into the freshwater. This leads to the decrease in concentration of these electrolytes in the organism's internal environment[16-18,20,22] (Figure 2).

Decline of the level of sodium and chlorine ions in the blood plasma of freshwater fish that takes place during the initial stage of stress is accompanied by drop of the osmolality of the organism internal environment[22]. As a result the osmotic pressures in the extracellular and intracellular liquids of the organism differ sharply and this jump of pressures facilitates transfer of water into the cells leading to the swelling of the latter. At hyponatremia the water



**Figure 2.** Dynamics of Na, K, concentrations in plasma of the roach at acute stress[16,18], and during different seasons[3,4].

content in the muscles of roach increases by 1.8%[23]. The increase in the water content in brain is especially injurious. The skull prevents expansion of the brain tissue. This is the brain edema at hyponatremia is accompanied by the rise of intracranial pressure with a range of relative negative consequences including high mortality[24].

It was found that during initial two to four hours of stress the leakage of potassium ions from cells prevail resulting in sharp increase in potassium concentration in the blood plasma[16,25] (Figure 2). Considerable increase in the blood plasma potassium content and its decrease in the cells lead to the decline in the gradient of this cation between tissue cells and intracellular liquid. This results in the drop of the membrane potential of cells. The membrane potential is a source of physical energy used for various functions of cells including secondary transport of a range of substances via surfaces of cellular membranes[26]. Its decrease at stress should lead to the decrease in the ability to fulfill transport functions. Many enzymes are situated on the cellular membranes at their activities depend on the value of cell membrane potential. It was revealed that during the initial stage of stress in common carp the level of potassium in the extracellular liquid increases[27]. This should result in the decrease in the membrane potential and is accompanied by inhibition of acetylcholine esterase activity.

To withstand damaging processes at stress (Figure 1, right part) the protective functions are enhanced by nervous and endocrine systems. These functions relate to the GAS (Figure 1, left part).

## 2.2. Protective functions at stress

The stressors perceived by nervous system serve as “alarm signals” entering various sectors of brain and hypothalamus. Hypothalamus modifies and coordinates the perceived information and affects the organism via various neuroendocrine pathways.

### 2.2.1. Hypothalamic-sympathetic-chromaffin cell axis

The first pathway relates to the release of catecholamines from the chromaffin cells, mainly via the preganglion cholinergic filaments of the sympathetic nervous system. However, the transection of these nerves does not prevent fully the secretion of catecholamines during the stress. It is suggested that in fish other factors (*e.g.* transported by the blood) participate in the above mentioned process[28]. Activation of chromaffin system at stress[29,30] leads to fast and sharp (by tenths times) concentration of catecholamines in blood plasma [28,30-34]. These hormones are responsible for various functions. They accelerate the consumption of oxygen by gill by means of increase in the rate of respiration, gill blood current, diffusion capacity of gill and transporting capacity of blood for oxygen via  $\beta$ -adrenergic receptors[28].

Catecholamines enhance the energetic metabolism. These hormones cause disintegration of polymers to monomers owing to increase in glycogenolysis, lipolysis and gluconeogenesis[35-38]. In addition, these hormones enhance oxidative function of mitochondria, functioning of the respiratory chain, oxygen consumption and, as a result, the generation of energy in a form of ATP[39].

### 2.2.2. Hypothalamic-pituitary-interrenal axis

The second pathway of the realization of protective processes at stress relates to the hypothalamic-pituitary-interrenal axis typical for terrestrial vertebrates and fish[40,41]. Hypothalamus secretes corticotrophin-releasing factor, arginine-vasotocin, isotocin and neuropeptide Y at stress. These substances reach hypophysis and stimulate releasing of adrenocorticotrophic hormone, melanophor-stimulating hormone (MSH) and beta-endorphin[42-44] (Figure 1, left part). As a result, the concentration of these hormones in the blood plasma increases[45-47]. The blood current delivers these hormones into the interrenal tissues enhancing the activity of the latter[29], and increase in the concentrations of cortisol[48-53], and corticosterone in the blood plasma[54].

Cortisol is the main glucocorticoid hormone in fish. It provides various functions: inhibits synthesis of proteins and enhances their degradation in the lymphoid, connective and muscle tissues. The amino acids released from proteins metabolize fast resulting in the negative nitrogen balance. These catabolic processes are determined by the induction of enzymes, especially of the enzymes responsible for catabolism of amino acids and for gluconeogenesis[39,55,56]. As a result, the synthesis of proteins is suppressed which leads to the growth rate retardation. In addition, cortisol stimulates metabolism of lipids by decomposition of fat. As a result, the level in fatty acids increases in the blood plasma[57,58]. Treatment of cortisol leads to the decrease in the level of lipids in the liver and muscles of coho salmon[57].

The reactions related to the energetic metabolism include the antioxidant system. The action of this system is directed towards the neutralization of the free radical processes[8]. The activation of energetic metabolism at stress is "automatically" accompanied by the enhancement of the antioxidant function. Catecholamines and corticosteroids stimulate the energetic metabolism and enhance the antioxidant system.

It is known that during the initial stage of stress the resistance of the organism decrease (the "alarm reaction")[59]. This indicates that at that time the effects of damage and malfunctions prevail over the adaptive reactions. It is suggests that at stress high levels of circulating catecholamines and corticosteroids in the blood plasma cause various disorders, including hyperhydration, loss of electrolytes and nonelectrolytes[32,33,41,60-62] as well as decrease in the resistance of erythrocytes to the blood current[63] in the freshwater fish. It is likely that this opinion is based upon the formal correlation between the increase in the concentrations of various hormones in the blood plasma and simultaneous manifestation of the damaging processes noted at stress. However, in reality the situation is different. Hypophysectomy[64], interrenalectomy and stanniusectomy[65] carried out separately and resulted in the desalination of the fish in the freshwater. The same phenomena are registered at the initial stage of stress in the intact animals resembling the situation when the endocrine glands do not function or excise. The situation is paradoxical: on the one hand at stress the functions of endocrine systems enhance accompanied by the increase in the concentrations of various hormones in the blood plasma, but on the other hand, the disorder of water-salt metabolism takes place at the high functional activity of the endocrine systems (Figure 2) which is typical at the excision of endocrine organs.

This phenomenon indicates at the initial stage of acute stress the hormones are seemingly deficient. At the very beginning of the stress response the protective functions related to the GAS (Figure 1, left part) do not compensate for damages (Figure 1, right part). As a result, the latter processes prevail resulting in the decrease of the organism resistance (the "alarm reaction" according to Selye[59]). The protective functions can not withstand development of damaging consequences of the stress instantly. Only with time the adaptive mechanisms repair the damages recovering and stabilizing the organism parameters either at new or at initial levels (Figure 2). This provides an improved stability for an organism (the "stage of resistance" according to Selye[59]). Long and strong stressor causes the exhaustion of the adaptive resources in the organism leading to the prevalence of destabilizing symptoms over the protective reactions (the "stage of exhaustion" according to Selye[59]).

### 3. Dynamics of physiological-biochemical parameters during the reproduction period in freshwater fish

As at stress the fish during the pre-spawning period exhibit enhancement of protective functions related to GAS (Figure 1, left part). In various fishes shortly before the spawning the activity of hypothalamus hypophysis neurosecretory systems increases which is accompanied by secretion of neurohormones of preoptic-hypophyseal neurosecretory system to blood[1,2]. With blood current the hormones of hypophysis reach interrenal gland[66-69] stimulating release of various corticosteroids into blood[66-71] including cortisol[72-80].

Corticosteroids realize transition of metabolism to the catabolic pathway owing to acceleration of gluconeogenesis, process of formation of glucose from noncarbohydrate sources[55] including protein. Concentration of proteins and amino acids decreases before spawning in fish blood serum and tissues[81-84]. Enhancement of gluconeogenesis and dissipation of glycogen (mainly in the liver) result in the increase blood plasma glucose which takes place before spawning as well[85]. Further the glucose is used for the energetic purposes. In addition, during the pre-spawning period the amount of accumulated fat in the fish body and various organs and tissues sharply decreases[83,86-89]. This indicates the increase in the share of lipolysis in the energetic metabolism.

Before spawning the activation of sympathetic nervous system accompanied by the increase in the blood catecholamine concentrations takes place in fish[90]. Catecholamines acting via  $\beta$ -adrenergic receptors accelerate consumption of oxygen by fish gill increasing the rates of respiration, blood current in gill, diffusion capacity of gill and transporting capacity of blood for oxygen[28].

It was revealed that before spawning the oxygen consumption rise in the whole organism and in separate tissues rises[91,92]. In addition, catecholamines enhance gluconeogenesis[36] and aerobic metabolism by increase in the Krebs cycle substrates[93]. It was also shown that the activity of the enzymes involved in Krebs cycle increases before spawning[94,95].

As it was shown above during the initial stage of at the development of GAS (Figure 1, left part) the signs of malfunctions of ionic homeostasis were observed (Figure 2). At the pre-spawning

period the situation is opposite: at the increase in the manifestations of GAS the protective functions related to maintaining of the ionic homeostasis prevail. For instance, before spawning in the roach blood plasma the concentration of sodium significantly increased by 10.1%; of potassium, decreased by 6.2 times (Figure 2). Pre-spawning increase in the content of sodium was observed in fish species as well[96,97]. This relates to the increase in the concentrations of various hormones in the blood plasma.

It was shown that cortisol enhances uptake of sodium and chlorine ions in the fresh-water fish gill from the environment[65,98]. This hormone facilitates differentiation of chloride cells in gill and increase specific activity of ion-transporting enzymes in gill, intestine and kidney[99,100]. Injections of cortisol and prolactin to hypophysectomized eel *Anguilla anguilla* L. increased concentration of sodium in the blood plasma to normal values and facilitated dehydration of tissues[101]. Adrenalin enhances uptake of sodium by gill[102] and prevents desalination of the organism[103] in the fish adapted to freshwater. The level of prolactin in the fish blood plasma rises before spawning[104]. This hormone decreases elimination of sodium from the organism by decreases in the permeability of gill epithelium and increased reabsorption of this ion in kidney[105].

Before spawning the level of potassium decreases considerably in the blood plasma of roach (Figure 2) and other fish species[97,106,107]. The fish blood plasma potassium level decrease in pre-spawning may relate to a certain extent to the effect of catecholamines. The experiments *in vitro* revealed that these hormones accelerated transport of potassium ions from plasma to erythrocytes of fish[108-112], frogs and pigeons[113]. Considerable decline of potassium level in blood causes increase in the gradient of concentrations of this cation between tissue cells and extracellular fluid. This leads to growth of the membrane potential in the cells. The membrane potential is a source of physical energy used for various functions of cell[26] including increase in the membrane-bound enzymes. For example, considerable increase in the activity of acetylcholine esterase and content of soluble fractions of proteins was revealed in the roach brain before spawning[114]. Increase in the physiological activity of the fish organism before spawning is accompanied by increase in the use of both forms of energy: chemical (ATP) and physical (membrane potential).

Comparative analysis revealed that in the freshwater fish with nonrecurring spawning the protective functions related to GAS enhance both at the initial period of stress and during the pre-spawning period (Figure 1, left part). As for the damaging processes (Figure 1, right part) the pattern of their dynamics during the initial stage of stress is strikingly different: at the initial stage of stress the damaging effects dominate over the protective processes which results in the decrease in the resistance of the organism. As opposite, during the pre-spawning period at the increase in GAS the protective functions are prevailing facilitating increase in the resistance of spawner. It was shown that during the pre-spawning period the resistance to stressors increase[97] along with increase in the thermal resistance in muscles[115] while the concentration of calcium in the erythrocytes and muscle tissues decrease[3,116]. Reduction of the intracellular calcium leads to the stabilization of cytoskeleton[117], determining increase in the resistance to any damaging factors at the level of cell.

The facts given above indicate that during the initial stage of stress due to domination of the damaging effects over the protective reactions the level of resistance of an organism decreases which corresponds to the Selye's "alarm reaction"[59]. Before spawning only protective systems dominate facilitating increase in the level of resistance in the spawner (Selye's "resistance stage"). That is, the absence of the alarm reaction is a characteristic of the pre-spawning period.

Why during the initial stage of stress at the enhancement of adaptive functions (Figure 1, left part) the damaging processes are manifested (Figure 1, right part) while, as opposite, before spawning only protective reactions prevail? This is because the stressors fall upon the organism suddenly enhancing simultaneously both the protective functions and the processes related to the damage of the affected organism. At this situation the protective system activated in the response to stress impact are unable to instantly withstand the damaging effects of extreme factors. As a result, for some time (the durations of this period depends on the strength of the factors and extent of the damage) the damaging processes dominate over the adaptive reactions causing decrease in the organism resistance (the "alarm reaction"). With time, strengthen protective functions eliminate stress-induced malfunctions and provide higher resistance to the organism (the "resistance stage").

As opposite, there are no sudden impacts before the spawning. This is why there are no damages to the organism. The state resembling strong stress occurs in the spawner only at the moment of spawning particularly (Figure 2). Both during the initial stage of stress and during the spawning the content of blood plasma sodium in fish drops sharply while the content of potassium rises. At that, in terms of the strength of the effect the moment of spawning is comparable to the acute stress (Figure 2). This means that during the stress the physiological and biochemical systems of spawner are subject to the strain comparable to that one at the extreme impacts. After the spawning, the resistance of fish decreases[97, 115,118] indicating the exhaustion of the efficiency of protective functions. This state corresponds to the Selye's "stage of exhaustion"[59]. The analysis of the muscles of roach immediately after the spawning confirms the exhaustion of the active physiological systems. During this period the concentration of sodium in the muscles rose by 16.9 mmol/kg wet mass; of potassium, as opposite, decreased by 27.5 mmol/kg wet mass[4]. Presumably, these changes take place owing to insufficient ability (exhaustion) of the sodium pump in the cellular membranes to evacuate the sodium ions in exchange for potassium. Then, within two to three weeks, both at stress and after spawning the parameters of ionic homeostasis recover and stabilize within the normal limits (Figure 2).

This clarifies the biological meaning of the enhancement of protective functions (Figure 1, left part) prior to the beginning of the spawning-induced impact. In the series of generations the acts of spawning repeat in a certain fish species at similar ecological conditions. Such cyclically standard situation promoted a set of adaptations triggering the protective systems in advance to the stress load related to spawning and forming a sort of "safety factor". If before the spawning such safety factor was not developed the reproducing fish may have died either during the spawning or after it owing to sharp decrease in the resistance and in the ability to maintain the ionic homeostasis below the permissible

levels. The post-spawning mortality in some less resistant fish is documented[119].

As it was noted above it is believed that during the stress high levels of catecholamines and corticosteroids in the blood plasma cause various malfunctions including impairment of the water-salt homeostasis[32,33,41,60-62,120] in freshwater fish. In this respect the studies on various functions during the pre-spawning period is an ideal model for revealing real effects of neuroendocrine system. During this period the neural and endocrinal functions enhance long before the spawning induced stress. Since before spawning no extreme factor affects the organism there are no damaging effects acting. This is why before spawning the effects of neural and endocrinal systems are manifested per se without simultaneous effects of spawning-related damaging factors.

No damaging processes affect fish during the pre-spawning period. As opposite during this period the protective effects dominate resulting in the increases in the organism resistance. If neuroendocrine systems causes damages to organism as it is supposed by many researchers[32,33,40,41,60-62,120], then the malfunctions should be evident during the pre-spawning period. However it is not occur. Hence, the enhancement of the neural and endocrinal systems during the initial stage of stress and before spawning is directed towards the providing of only the adaptive functions.

#### 4. Conclusions

The comparative analysis revealed that the enhancement of protective functions related to the GAS takes place in the reproducing freshwater fish with nonrecurrent spawning both at the initial stage of stress and during the pre-spawning period. However, during the initial stage of stress the damaging effects dominate over the adaptive processes resulting in the decrease in the organism resistance (the “alarm reaction” according to Selye[59]). As opposite, during the pre-spawning period at the intensification of GAS only the protective systems dominate facilitating the increase in the level of resistance in the spawner (the “stage of resistance” according to Selye[59]). The alarm stage is not manifested before spawning. The state comparable to strong stress occurs in the reproducing fish only at the act of spawning. This is accompanied by decrease in the organism resistance. This indicates the exhaustion of the protective functions and corresponds to the Selye’s “stage of exhaustion”[59]. With time (within two to three weeks), both at stress and after spawning the physiological-biochemical parameters recover and stabilize within the normal limits.

#### Conflict of interest statement

I declare that I have no conflict of interest.

#### Acknowledgments

This study has been financially supported by the budgetary state project: Studying of adaptations of organisms, populations, communities of aquatic organisms and modeling ecosystems to environmental factors. State registration No. 01201351909.

#### References

- [1] Garlov PE. [Stress as a state of “species-specific” physiological normal simultaneous spawning in some fish species]. In: Ozernyuk ND, editor. *[Ecological problems of fish ontogenesis]*. Moscow: Moscow State University Publ; 2001, p. 266-282. Russian.
- [2] Garlov PE, Polenov AL. [Functional cytology of preoptical-pituitary neurosecretory system in fish]. *Tsitologiya* 1996; **38**: 275-299. Russian.
- [3] Martemyanov VI. [Ranges of regulation of sodium, potassium, calcium, magnesium concentrations in plasma, erythrocytes, and muscle tissue of *Rutilus rutilus* under natural conditions]. *J Evol Biochem Physiol* 2001; **37**: 141-147. Russian
- [4] Martemyanov VI. [Dynamics of content of cations in plasma, erythrocytes and muscle tissues of roach, *Rutilus rutilus* L. during the reproductive period]. *Biol Inland Waters* 2004; **2**: 78-84. Russian.
- [5] Martemyanov VI. [Stress in fish: protective and damaging processes]. *Biol Inland Waters* 2002; **4**: 3-13. Russian.
- [6] Mallatt J. Fish gill structural changes induced by toxicants and other irritants: a statistical review. *Can J Fish AquatSci* 1985; **42**: 630-648.
- [7] Brekhman II, Golotin VG, Gonenko VA, Dobryakova AI. [On the accumulation of lipid peroxides and quinines in the liver or rats at stress]. In: Danilov VS, editor. *[Bioantioxidants and regulation of oxidative processes in cell]*. Moscow: Moscow State University Publ; 1972, p. 18-23. Russian.
- [8] Voskresensky ON. [On the relations of adaptogenic and antioxidant effects]. In: Brekhman II, editor. *[Adaptation and adaptogenesis]*. Vladivostok: DVNTS AN SSSR; 1977, p. 91-96. Russian.
- [9] Golotin VG. [Toxic peroxides at stress]. In: Brekhman II, editor. *[Adaptation and adaptogenesis]*. Vladivostok: DVNTS AN SSSR; 1977, p. 33-35. Russian.
- [10] Kolosova NG, Shorin YP, Kulikov V Yu. [The reactions of lipid peroxide oxidation in liver and lung of rats under longterm cold adaptation]. *Byull Eksp Biol Med* 1981; **4**: 436-437. Russian.
- [11] Chernyshev VI. [Etiology and prevention of free radical pathology during physiological adaptation of fish to unusual ecological conditions]. *Trudy Moskovskogo Obschestva Ispytatelei Prirody, Otd Biol* 1982; **57**: 141-150. Russian.
- [12] Chernyshev VI, Isuev AR. [Etiology of free radical pathology in Acipenseridae *Acipenser guldenstadti* Brant, *Acipenserstellatus pallas*, *Huso huso* (L.) during embryogenesis]. *Vopr Ikhtiol* 1980; **20**: 334-344. Russian.
- [13] Chernyshev VI, Telitchenko MM. [Physico-chemical aspects of development of pathological processes in *Cyprinus carpio* L. at sharp changes of temperature, water concentration oxygen, and intoxication with poisons of different nature]. *Vopr Ikhtiol* 1973; **13**: 155-166. Russian.
- [14] Geraskin PP, Ponomareva EN, Metallov GF, Galaktionova ML. [Oil pollution of the Caspian Sea as one of the factors of initiation of oxidative stress in sturgeons]. *Izvestiy Samarskogo Nauchnogo Tsentra RAN* 2012; **14**: 209-216. Russian.
- [15] Magomedov SK, Chernyshev VI, Shekhanova IA. [Comparative study of some physico-chemical mechanisms of fish adaptation to different water salinity]. *Vopr Ikhtiol* 1974; **14**: 492-498. Russian.
- [16] Martemyanov VI. [Dynamics of electrolyte concentrations in fresh-water

- fish during stress]. In: Monakov AV, editor. *[Fresh-water fauna and biology of hydrobionts]*. Leningrad: Nauka; 1983, p. 237-248. Russian.
- [17] Martemyanov VI. [Based on ion exchange data methods for assessment of unfavorable environment exposure on aquatic organisms]. *Water: Chem Ecol* 2013; **3**: 52-63. Russian.
- [18] Martemyanov VI. Patterns of changes in sodium content in plasma and erythrocytes of freshwater fish at stress. *J Ichthyol* 2013; **53**: 220-224.
- [19] Vinogradov GA, Kelrman AK. [Ion exchange in freshwater fishes under stress]. *Vopr Ikhtiol* 1987; **272**: 307-312. Russian.
- [20] Postlethwaite E, McDonald D. Mechanisms of Na<sup>+</sup> and Cl<sup>-</sup> regulation in freshwater-adapted rainbow trout (*Oncorhynchus mykiss*) during exercise and stress. *J Exp Biol* 1995; **198**: 295-304.
- [21] Meyer DK. Physiological adjustments in chloride balance of the goldfish. *Science* 1948; **108**: 305-307.
- [22] Natochin Yu V, Shakhmatova EI, Lavrova EA, Maksimovich AA, Karpenko LA, Martemyanov VI. [Volumoregulation in cells of freshwater and migrating fish during changes in osmolality and ionic composition of the blood serum]. *J Evol Biochem Physiol* 1991; **272**: 159-166. Russian.
- [23] Martemyanov VI. [Hyponatremia in water and land vertebrates as criterion of adverse physiological state of organisms affected by adverse environmental factors]. In: Organisms, populations, ecosystems: problems and ways of biodiversity preservation, proceedings of all Russian Conference; 2008 Nov 24-28; Vologda, Russian. 2008. p. 67-71. Russian.
- [24] Arief AI. Hyponatremia, convulsions, respiratory arrest, and permanent brain damage after elective surgery in healthy women. *N Engl J Med* 1986; **314**: 1529-1535.
- [25] Martemyanov VI, Zaprudnova RA. [Dynamics of electrolyte concentration in the blood plasma, red blood cells, and muscle tissue of fresh-water fish during stress]. *Biol Nauki* 1982; **10**: 44-49. Russian.
- [26] Skulachev VP. *[Energetics of biological membranes]*. Moscow: Nauka; 1989. Russian.
- [27] Kozlovskaya VI, Martemyanov VI. Brain acetyl cholinesterase activity in carp (*Cyprinus carpio* L.) during acute and chronic phenol intoxication. *Hydrobiol J* 1991; **27**: 75-81.
- [28] Randall DJ, Perry SF. Catecholamines. In: Hoar WS, Randall DJ, Farrell AP, editors. *Fish physiology*. Vol. 12B. San Diego: Academic Press; 1992, p. 255-300.
- [29] Mezhnin FI. [Interrenal and suprarenal glands and *Stannius corpuscles* in guppy, *Lebistes reticulatus* Peters in the conditions of extreme stress]. *Vopr Ikhtiol* 1978; **18**: 697-718. Russian.
- [30] Zaprudnova RA, Prozorovskaya MP [The change in concentrations of catecholamines and ions in tissues of bream (*Abramis brama*) under stress]. *J Ichthyol* 1999; **39**: 262-266.
- [31] Gingerich WH, Drott KR. Plasma catecholamine concentrations in rainbow trout (*Salmo gairdneri*) at rest and after anesthesia and surgery. *Gen Comp Endocrinol* 1989; **73**: 390-397.
- [32] Mazeaud MM, Mazeaud F. Adrenergic responses to stress in fish. In: Pickering AD, editor. *Stress and fish*. London: Academic Press; 1981, p. 49-75.
- [33] Mazeaud MM, Mazeaud F, Donaldson EM. Primary and secondary effects of stress in fish: some new data with a general review. *Trans Am Fish Soc* 1977; **106**: 201-212.
- [34] McDonald DG, Milligan CL. Chemical properties of the blood. In: Hoar WS, Randall DJ, Farrell AP, editors. *Fish physiology*. Vol. 12B. San Diego: Academic Press; 1992, p. 55-133.
- [35] Danulat E, Mommsen TP. Norepinephrine: a potent activator of glycogenolysis and gluconeogenesis in rockfish hepatocyte. *Gen Comp Endocrinol* 1990; **78**: 12-22.
- [36] Mommsen TP, Walsh PJ, Perry SF, Moon TW. Interactive effects of catecholamines and hypercapnia on glucose production in isolated trout hepatocytes. *Gen Comp Endocrinol* 1988; **70**: 63-73.
- [37] Wright PA, Perry SF, Moon TW. Regulation of hepatic gluconeogenesis and glycogenolysis by catecholamines in rainbow trout during environmental hypoxia. *J Exp Biol* 1989; **147**: 169-188.
- [38] Pickering AD, Pottinger TG. Biochemical effects of stress. In: Hochachka PW, Mommsen TP, editors. *Environmental and ecological biochemistry*. Amsterdam: Elsevier; 1995, p. 349-379.
- [39] Kulinsky VI, Olkhovsky IA. [Two adaptation strategies under unfavorable conditions: the resistant and tolerant ones role of hormones and receptors]. *Usp Sovrem Biol* 1992; **112**: 697-714. Russian.
- [40] Gronov G. [Application of concept a "Stress" for bony fishes]. *Zool Anz* 1974; **192**: 316-331. German.
- [41] Wendelaar Bonga SE. The stress response in fish. *Physiol Rev* 1997; **77**: 591-625.
- [42] Lederis K, Fryer JN, Okawara Y, Schönrock C, Richter D. Corticotropin releasing factors acting on the fish pituitary: experimental and molecular analysis. In: Hoar WS, Randall DJ, Farrell AP, editors. *Fish physiology*. Vol. 13. San Diego, Academic Press; 1994, p. 68-100.
- [43] Olivereau M, Olivereau JM. Corticotropin-like immunoreactivity in the brain and pituitary of three teleost species (goldfish, trout and eel). *Cell Tissue Res* 1990; **262**: 115-123.
- [44] Olivereau M, Olivereau J. Responses of brain and pituitary immunoreactive corticotropin-releasing factor in surgically interrenalectomised eels: immunocytochemical study. *Gen Comp Endocrinol* 1991; **81**: 295-303.
- [45] Balm PH, Pottinger TG. Corticotrope and melanotrope POMC-derived peptides in relation to interrenal function during stress in rainbow trout (*Oncorhynchus mykiss*). *Gen Comp Endocrinol* 1995; **98**: 279-288.
- [46] Pickering AD, Pottinger TG, Sumpter JP. Independence of the pituitary-interrenal axis and melanotroph activity in the brown trout *Salmo trutta* L. under conditions of environmental stress. *Gen Comp Endocrinol* 1986; **64**: 206-211.
- [47] Sumpter JP, Dye HM, Benfey TJ. The effects of stress on plasma ACTH, alpha-MSH, and cortisol levels in salmonid fishes. *Gen Comp Endocrinol* 1986; **62**: 377-385.
- [48] Abreu JS, Takahashi LS, Hoshiiba MA, Urbinati EC. Biological indicators of stress in pacu (*Piaractus mesopotamicus*) after capture. *Braz J Biol* 2009; **69**: 415-421.
- [49] Barton BA, Iwama GK. Physiological changes in fish from stress in aquaculture with emphasis on the responses and effects of corticosteroids. *Ann Rev Fish Dis* 1991; **1**: 3-26.
- [50] Barton BA. Stress in fishes: a diversity of responses with particular reference to changes in circulating corticosteroids. *Integr Comp Biol* 2002; **42**: 517-525.
- [51] Carragher JF, Rees CM. Primary and secondary stress responses in golden perch, *Macquaria ambigua*. *Comp Biochem Physiol A* 1994; **107**: 49-56.
- [52] Donaldson EM. The pituitary-interrenal axis as an indicator of stress in

- fish. In: Pickering AD, editor. *Stress in fish*. London: Academic Press; 1981, p. 11-47.
- [53] Pickering AD, Pottinger TG. Stress responses and disease resistance in salmonid fish: effects of chronic elevation on plasma cortisol. *Fish Physiol Biochem* 1989; **7**: 253-258.
- [54] Martemyanov VI. [The dynamics of corticosterone and electrolytes concentrations in the bream blood plasma under stress conditions]. *Biol Vnutrennikh Vod Informatsionnyi Byull* 1987; **75**: 51-54. Russian.
- [55] Janssen PA, Waterman J. Hormonal regulation of gluconeogenesis and glycogenolysis in carp (*Cyprinus carpio*) liver pieces cultured *in vitro*. *Comp Biochem Physiol A* 1988; **91**: 451-455.
- [56] Vijayan MM, Reddy PK, Leatherland JF, Moon TW. The effects of cortisol on hepatocyte metabolism in rainbow trout: a study using the steroid analogue RU486. *Gen Comp Endocrinol* 1994; **96**: 75-84.
- [57] Sheridan MA. Lipid dynamics of fish: aspects of absorption, transportation, deposition and mobilization. *Comp Biochem Physiol B* 1988; **90**: 679-690.
- [58] Sheridan MA. Regulation of lipid metabolism in poikilothermic vertebrates. *Comp Biochem Physiol B* 1994; **107**: 495-508.
- [59] Selye G. [At the level of the whole organism]. Moscow: Nauka; 1972. Russian.
- [60] Gonzalez R, McDonald D. The relationship between oxygen consumption and ion loss in a freshwater fish. *J Exp Biol* 1992; **163**: 317-332.
- [61] McDonald DG, Cavdek V, Ellis R. Gill design in freshwater fishes interrelationships among gas exchange, ion regulation, and acid-base regulation. *Physiol Zool* 1991; **64**: 103-123.
- [62] Nilsson S. Control of gill blood flow. In: Nilsson S, Holmgren S, editors. *Fish physiology: recent advanced*. London: Croom Helm; 1986, p. 87-101.
- [63] Wells RMG, Davie PS, Weber RE. Effect of beta-adrenergic stimulation of trout erythrocytes on blood viscosity. *Comp Biochem Physiol C Comp Pharmacol* 1991; **100**: 653-655.
- [64] Ensor DM, Ball JN. Prolactin and osmoregulation in fishes. *Fed Proc* 1972; **31**: 1615-1623.
- [65] Chester Jones I, Chan DKO, Henderson IW, Ball JN. The adrenocortical steroids, adrenocorticotropin and the corpuscle of stannius. In: Hoar WS, Randall DJ, editors. *Fish physiology*. Vol. 2. London: Academic Press; 1969, p. 322-376.
- [66] Barannikova IA, Vasilyeva EV, Trenkler IV, Tsepelovan PG. [The interregal gland in the life cycle of anadromous sturgeons (fam. Acipenseridae)]. *Vopr Ikhtiol* 1978; **18**: 719-734. Russian.
- [67] Heyl HL, Carpenter SJ. Reversible changes in adrenal cortical cell morphology and plasma hydroxycorticosteroids during freshwater portion of the spawning journey of Atlantic salmon (*Salmo salar*). *J Fish Res Board Can* 1972; **29**: 311-314.
- [68] Pen'kova EA. [On the functional changes in the interregal tissue of adult sturgeon *Acipenser guldenstadti* Brandt during the life cycle]. *Vopr Ikhtiol* 1974; **14**: 134-143. Russian.
- [69] Vasilyeva EV, Barannikova IA. [Ultrastructure of the cells of interregal tissue of sturgeon and its comparative analysis before and after spawning]. *Tsitologiya* 1978; **20**: 263-268.
- [70] Hane S, Robertson OH. Changes in plasma 17-hydroxycorticosteroids accompanying sexual maturation and spawning of the Pacific salmon (*Oncorhynchus tshawytscha*) and rainbow trout (*Salmo gairdnerii*). *Proc Natl Acad Sci U S A* 1959; **45**: 886-893.
- [71] Tsepelovan PG, Rusakov Yu I. [Corticosteroid content of the blood in the sturgeon *Acipenser guldenstadti* at various stages of life cycle]. *Zh Evol Biokhim Fiziol* 1976; **12**: 77-80. Russian.
- [72] Bry C. Plasma cortisol levels of female rainbow trout (*Salmo gairdneri*) at the end of the reproductive cycle: relationship with oocyte stages. *Gen Comp Endocrinol* 1985; **57**: 47-52.
- [73] Fuller JD, Scott DBC, Fraser R. Effects of catching techniques, captivity and reproductive cycle on plasma cortisol concentration in the powan (*Coregonus lavaretus*), a freshwater teleost from Loch Lomond. *J Endocrinol* 1975; **63**: 1-24.
- [74] Hou Y, Han X, Suzuki Y. Annual changes in plasma levels of cortisol and sex steroid hormones in male rainbow trout, *Oncorhynchus mykiss*. *Chin J Oceanol Limnol* 2001; **19**: 217-221.
- [75] Kusakabe M, Nakamura I, Young G. 11beta-hydroxysteroid dehydrogenase complementary deoxyribonucleic acid in rainbow trout: cloning, sites of expression, and seasonal changes in gonads. *Endocrinology* 2003; **144**: 2534-2545.
- [76] Noaksson E, Linderöth M, Gustavsson B, Zebuhr Y, Balk L. Reproductive status in female perch (*Perca fluviatilis*) outside a sewage treatment plant processing leachate from a refuse dump. *Sci Total Environ* 2005; **340**: 97-112.
- [77] Pickering AD, Christie P. Changes in the concentrations of plasma cortisol and thyroxine during sexual maturation of the hatchery-reared brown trout, *Salmo trutta* L. *Gen Comp Endocrinol* 1981; **44**: 487-496.
- [78] Pickering AD, Pottinger TG. Lymphocytopenia and interrenal activity during sexual maturation in the brown trout, *Salmo trutta* L. *J Fish Biol* 1987; **30**: 41-50.
- [79] Westring CG, Ando H, Kitahashi T, Bhandari RK, Ueda H, Urano A, et al. Seasonal changes in CRF-I and urotensin I transcript levels in masu salmon: correlation with cortisol secretion during spawning. *Gen Comp Endocrinol* 2008; **155**: 126-140.
- [80] Wingfield JC, Grimm AS. Seasonal changes in plasma cortisol, testosterone and oestradiol-17β in the plaice, *Pleuronectes platessa* L. *Gen Comp Endocrinol* 1977; **31**: 1-11.
- [81] Dzhabarov MI. [Dynamics of contents of hemoglobin and total protein in the blood serum of common carp, bream and shemaya near the Kura River mouth area of the Black Sea in relation to maturation of gonads]. In: Grese VN, editor. *Energetic aspects of growth and metabolism in aquatic animals*. Kiev: Naukova Dumka; 1972, p. 51-52. Russian.
- [82] Fedorova LS, Grudanova SD. [Some features of protein metabolism in the females beluga sturgeon during spring migration and maturation of gonads]. In: *Development of biological bases and biotechnology of sturgeon fisheries in the USSR waterbodies (on the Basis of 1967 Materials)*. Astrakhan; TsNIORKhpubl; 1968, p. 73-76. Russian.
- [83] Golovanenko LF, Shuvatova TF, Putina EP, Fedorova LS, Arakelova AA. [Physiological-biochemical characteristics of the Don River zander at various stages of sexual cycle]. *Vopr Ikhtiol* 1970; **10**: 374-384. Russian.
- [84] Naidenova IN. [Changes in the content of protein in the organs of horse mackerel and scorpion fish during the process of gonads development]. In: Karzinkin GS, editor. [Metabolism and biochemistry of fishes]. Moscow: Nauka; 1967, p. 314-316. Russian.
- [85] Nace PF, Moule ML, Schuh JE. The normal blood sugar of the toadfish. *Can J Physiol Pharmacol* 1964; **42**: 225-232.
- [86] Chugunova NI, Assman AV, Makarova NP. [Growth and dynamics of fatness in fish as adaptive processes]. In: Nikolsky GV, editor. *Problems*

- of fish ecology. Moscow: USSR Acad Sci Publ; 1961, p. 96-181. Russian.
- [87] Shatunovskii MI. [Dynamics of fat and water content of meat and gonads of the European flounder (*Pleuronectes flesus* Linne) and its relationship with the characteristics of gonad maturation]. *Vopr Ikhtiologii* 1963; **3**: 652-667. Russian.
- [88] Shikhshabekov MM. [Specificity of sexual cycles in some anadromous fish in the lower reaches of the Terek River]. *Vopr Ikhtiologii* 1974; **14**: 95-104. Russian.
- [89] Vyatchanin LI. [Dynamics of fat accumulation in relation to maturation of gonads in the roach of the Kremenchug Reservoir]. *Gidrobiologicheskii Zhurnal* 1971; **7**: 94-98. Russian.
- [90] Pustovoitova-Vasilene ME. [Seasonal changes in content of catecholamines and DOPA in tissues of the carp *Cyprinus carpio*]. *Zh Evol Biokhim Fiziol* 1972; **8**: 93-94. Russian.
- [91] Stroganov NS. [*Ecological physiology of fish*]. Moscow: Izdvo AN SSSR; 1962. Russian.
- [92] Yurkevich GN. [Seasonal changeability of tissue respiration in rock perches]. *Vopr Ikhtiologii* 1972; **12**: 202-204. Russian.
- [93] Kondrashova MN. [The scheme of deviations of state of mitochondria from the norm and the substances inducing these deviations]. In: Kondrashova MN, editor. *Reaction of live systems and state of energetic metabolism*. Pushchino: ONTINTsBI; 1979, p. 185-191. Russian.
- [94] Marquez ED. A comparison of glutamic-oxalacetate transaminase, lactate dehydrogenase, L-hydroxybutyrate dehydrogenase and creatine phosphokinase activities in non-spawning, pre-spawning and spawning pink salmon. *Comp Biochem Physiol B* 1976; **54**: 121-123.
- [95] Pye VI, Wieser W, Zech M. The effect of season and experimental temperature on the rates of oxidative phosphorylation of liver and muscle mitochondria from the tench *Tinca tinca*. *Comp Biochem Physiol B* 1976; **54**: 13-20.
- [96] Boiko P Ya, Maksimovich AA, Natochin Yu V, Polyakov VN, Plusnin VV. [Osmolality and electrolytes of blood serum in spawning gorbusha *Onchorhynchus gorbusha* during migration from sea to river and the reverse migration to the sea]. *Vopr Ikhtiologii* 1988; **28**: 467-472. Russian.
- [97] Zaprudnova RA, Martemyanov VI. [Seasonal changes in cation concentration in blood plasma of freshwater fish]. *Vopr Ikhtiologii* 1988; **28**: 671-676. Russian.
- [98] Laurent P, Parry SF. Effects of cortisol on gill chloride cell morphology and ionic uptake in the freshwater trout, *Salmo gairdneri*. *Cell Tissue Res* 1990; **259**: 429-442.
- [99] Chester Jones I, Mosley W, Henderson IW, Garland HO. The interrenal gland in pisces. In: Chester Jones I, Henderson IW, editors. *General, comparative and clinical endocrinology of the adrenal cortex*. Vol. 3. London: Academic Press; 1980, p. 396-523.
- [100] McCormick SD. Hormonal control of gill  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase and chloride cell function. In: Wood CM, Shuttleworth TJ, editors. *Fish physiology*. Vol. 14. New York: Academic Press; 1995, p. 285-315.
- [101] Chan DKO, Chester Jones I, Mosley W. Pituitary and adrenocortical factors in the control of water and electrolyte composition of the freshwater European eel (*Anguilla anguilla* L.). *J Endocrinol* 1968; **42**: 91-98.
- [102] Richards BD, Fromm PO. Sodium uptake by isolated-perfused gills of rainbow trout (*Salmo gairdneri*). *Comp Biochem Physiol* 1970; **33**: 303-310.
- [103] Pic P, Mayer-Gostan N, Maetz J. Branchial effects of epinephrine in the seawater-adapted mullet. II.  $\text{Na}^+$  and  $\text{Cl}^-$  extrusion. *Am J Physiol* 1975; **228**: 441-447.
- [104] Singh SP, Singh TP. Prolactin level in relation to gonadotrophin concentration during different phases of annual reproductive cycle in the freshwater catfish, *Clarias batrachus* (Linn.). *Ann Endocrinol* 1980; **41**: 415-424.
- [105] Lahlou B. [Hormones in osmoregulation of fishes]. In: Ali MA, editor. *Environmental physiology of fishes*. New York: Springer US; 1980, p. 201-240. Russian.
- [106] Lavrova EA, Natochin Yu V. [Salt composition of environment and cation concentration in blood serum in fish of different systematic groups]. In: Stroganov NS, editor. [*Ecological physiology of fish*]. Moscow: Nauka; 1973, p. 151-153. Russian.
- [107] Sokolova MM, Bakhteeva VT. [On the role of kidneys in potassium balance in fish]. In: Stroganov NS, editor. [*Current problems of ecological physiology of fish*]. Moscow: Nauka; 1979, p. 214-217. Russian.
- [108] Bourne PK, Cossins AR. On the instability of  $\text{K}^+$  influx in erythrocytes of the rainbow trout, *Salmo gairdneri*, and the role of catecholamine hormones in maintaining *in vivo* influx activity. *J Exp Biol* 1982; **101**: 93-104.
- [109] Houston AH, McCulloch CAM, Keen J, Maddalena C, Edwards J. Rainbow trout red cells *in vitro*. *Comp Biochem Physiol A* 1985; **81**: 555-565.
- [110] Houston AH, Gray JD. Red cell magnesium in rainbow trout. *Can J Zool* 1988; **66**: 929-933.
- [111] Zaprudnova RA. [Effect of adrenaline on plasma potassium concentration at incubation of the bream blood]. *Inform Byull Biol Vnutrennikh Vod* 1983; **58**: 55-58. Russian.
- [112] Zaprudnova RA. [Effect of adrenaline on the content of potassium in the blood plasma of bream at stress and spawning]. *Biol Inland Waters* 2000; **4**: 133-140. Russian.
- [113] Orskov SL. Experiments on the influence of adrenaline and noradrenaline on the potassium absorption of red blood cells from pigeons and frogs. *Acta Physiol Scand* 1956; **37**: 299-306.
- [114] Chuiko GM, Zhelnin Y, Podgornaya VA. Seasonal fluctuations in brain acetylcholinesterase activity and soluble protein content in roach (*Rutilus rutilus* L.): a freshwater fish from Northwest Russia. *Comp Biochem Physiol C* 1997; **117**: 251-257.
- [115] Altukhov Yu V. [Seasonal changes in thermostability of isolated muscle tissue in the Black Sea horse mackerel]. *Tsitologiya* 1963; **5**: 241-243. Russian.
- [116] Martemyanov VI. [Seasonal dynamics of calcium content in the plasma, erythrocytes, muscles and gonads of roach, *Rutilus rutilus* L.]. *Biol Inland Waters* 1998; **2**: 73-79.
- [117] Boldyrev AA. [*Introduction to membrane biochemistry*]. Moscow: Vysshaya Shkola; 1986. Russian.
- [118] Mikryakov VR. [Topical problems of immunology]. In: Butorin NV, Poddubnyi AG, editors. *Theoretical aspects of fisheries studies in the reservoirs*. Leningrad: Nauka; 1977, p. 83-97. Russian.
- [119] Nielsen JR, Lambert G, Bastardie F, Sparholt H, Vinther M. Do Norway pout (*Trisopterus esmarkii*) die from spawning stress? Mortality of Norway pout in relation to growth, sexual maturity, and density in the North Sea, Skagerrak, and Kattegat. *ICES J Mar Sci* 2012; **69**: 197-207.
- [120] Palikova M, Svobodova Z. [Biological indicators of stress at fishes (Review)]. *Bull VURH Vod* 1995; **31**: 17-27. Czech.