

Journal of Coastal Life Medicine

journal homepage: www.jclmm.com



Document heading doi: 10.12980/JCLM.3.201514J67

©2015 by the Journal of Coastal Life Medicine. All rights reserved.

Phytochemical studies of various polarities leave crude extracts of Omani *Datura metel* L. and evaluation of their antimicrobial potential

Ali Saleh Hamed Al-Jafari, Mohammad Amzad Hossain*

School of Pharmacy, College of Pharmacy and Nursing, University of Nizwa, P. O. Box 33, Postal Code 616, Nizwa, Sultanate of Oman

PEER REVIEW

Peer reviewer

Dr. Abu Bakar Siddque, Assistant Professor, Department of Drug Analysis, University of Dhaka, Bangladesh.
E-mail: 90zadid@gmail.com

Comments

The present study on biochemical screening of various leaves crude extracts of *D. metel* is giving the valuable brief and scientific information about this plant.
Details on Page 933

ABSTRACT

Objective: To identify the chemical constituents and evaluate antimicrobial potential of various crude extracts from leaves of *Datura metel* grown in Oman.

Methods: The leaf samples were collected from the University of Nizwa and extracted with methanol by using Soxhlet extractor. The isolated crude extract was defatted with distilled water and extracted with solvents of different polarities including hexane, chloroform, ethyl acetate and butanol. Chemical compositions of the crude extracts were analyzed by gas chromatography-mass spectrometer and their antimicrobial potential was evaluated by agar disc diffusion method against one Gram positive bacteria *Staphylococcus aureus* and two Gram negative bacteria *Escherichia coli* and *Pseudomonas aeruginosa*.

Results: The crude extracts were composed of different organic compounds such as alkaloids, hydrocarbons, aromatic hydrocarbons, organic acids, terpenoids, vitamin etc. The methanol and its fractionated crude extracts showed antimicrobial potential with inhibition zone in the range of 0-13 mm.

Conclusions: The selective crude extract from the leaves of *Datura metel* could be used as natural antibiotics.

KEYWORDS

Datura metel, Soxhlet extractor, Crude extracts, Gas chromatography-mass spectrometry analysis, Antimicrobial activity

1. Introduction

Since ancient times, different parts of medicinal plants have been used to cure specific ailments. The raw materials of medicinal and aromatic plants, crude extracts and essential oils contain various active secondary metabolites like tannins, terpenoids, alkaloids, flavonoids, phenols, steroids and glycosides[1]. Crude extracts of all plants especially medicinal plants are potential sources of antimicrobial, antifungal and antibacterial agents[2-4]. The resistance of bacterial strains against antibiotics makes many scientists and researchers recently pay attention to the crude extracts and

biologically active pure compounds isolated from plant species used in herbal remedies[5-9].

Datura metel L. (*D. metel*) is one of the most important herbal plants used worldwide for the treatment of different diseases[10-12]. *D. metel* is a flowering plant belonging to family Solanaceae. The maximum height of this plant is about 1.5 m. Its leaves are completely different; the leaves are simple, alternate, dark green, broadly ovate, shallowly lobed and glabrous. The leaves are traditionally used for treatment of diarrhoea, gonorrhoea and bronchitis[6-11,13-15]. They are also used for the treatment of catarrh, epilepsy, insanity, hysteria, rheumatic pains, hemorrhoids, painful

*Corresponding author: Mohammad Amzad Hossain, School of Pharmacy, College of Pharmacy and Nursing, University of Nizwa, P. O. Box 33, Postal Code 616, Nizwa, Sultanate of Oman.

Tel: +968 99708496

Fax: +968 25446236

E-mail address: hossainabi@gmail.com

Foundation Project: Supported by Central Instrument Laboratory, College of Agriculture and Marine Sciences, Sultan Qaboos University, Sultanate of Oman (Grant No. 507/SOP/OB/1/2013).

Article history:

Received 9 Sep 2014

Received in revised form 11 Sep 2014

Accepted 20 Sep 2014

Available online 5 Oct 2014

menstruation, skin ulcers and wounds[16-19]. Recently, the paste of leaves has been used in the treatment of burns[16]. In Nigeria, this plant is traditionally used to cure diseases such as asthma, cough, convulsion and insanity[19]. The Nigeria nationals are also using the leaves and seeds as anesthetic, antispasmodic, antitussive, bronchodilator and hallucinogenic[18]. The main chemical constituents of *D. metel* reported in the literature are alkaloids[20-22]. The main target of the present work was to isolate and identify chemical constituents and to evaluate the antimicrobial potential of various crude extracts of the leaves of *D. metel* which was native to Sultanate of Oman.

2. Materials and methods

2.1. Chemicals

Methanol, hexane, chloroform, ethyl acetate, butanol and amoxicillin were from Sigma-Aldrich Company Limited, Germany. Dichloromethane and acetone were from British Drug Houses, UK. Filter paper discs were obtained from Whatmann, China. Bacterial strains, including *Staphylococcus aureus* ATCC 29213 (*S. aureus*), *Escherichia coli* ATCC 9637 (*E. coli*), and *Pseudomonas aeruginosa* ATCC 9027 (*P. aeruginosa*) were collected from Nizwa Hospital, Nizwa, Sultanate of Oman.

2.2. Plant materials

The leaf samples of *D. metel* were collected from the University of Nizwa campus. After collection the samples were transported to the Natural Product Laboratory and kept at room temperature for cleaning, drying and extracting. Then fresh leaf samples were separated from the ones affected by fungus and bacteria.

2.3. Preparation of plant material

The unaffected leaf samples of *D. metel* were washed with water to remove the dust and foreign particles. After washing, the leaf samples were dried at room temperature for 3 weeks and ground into fine powder.

2.4. Extraction

The leaf powder sample (85 g) was extracted with methanol (300 mL) by using Soxhlet extractor until complete extraction. The methanol was completely evaporated by using rotary evaporator to give methanol free crude extract. The crude extract (33.7 g) was defatted with water (200 mL) and fractionated by using hexane, chloroform, ethyl acetate and butanol to give hexane, chloroform, ethyl acetate, butanol and residual hydro alcoholic fractions.

2.5. Antimicrobial assay

All crude extracts from the leaves of *D. metel* were tested for their antimicrobial activity by using three bacterial strains such as *S. aureus*, *E. coli* and *P. aeruginosa* grown on nutrient agar plates by using disc diffusion technique. Four concentrations 0.25, 0.50, 1.00 and 2.00 mg/mL were prepared for each crude extracts by using dimethyl sulphoxide as a solvent. The positive control amoxicillin was also prepared with the same solvent. Filter paper discs of 5 mm

in diameter were impregnated with each concentration and placed on the agar plate which was inoculated with standard bacteria strains. All the sample plates were incubated at 37 °C for 24 h. The antimicrobial activity was evaluated by measuring the diameter of inhibition zone against the tasted bacteria. Each experiment was done in triplicate.

2.6. Gas chromatography-mass spectrometry (GC-MS) analysis

The different polarities of crude extracts of *D. metel* were analysed by using a Perkin Elmer GC (Model Perkin Elmer Clarus 500, USA) coupled with a Perkin Elmer Clarus 600 C MS. The special fused silica capillary column was used for the analysis of crude extracts with specification of 30 m×0.25 mm ID, film thickness 0.25 µm. For the detection, separation and ionization by an electron ionization system with ionization energy 70 eV was performed. Helium was used as a carrier gas at constant flow rate of 1 mL/min. The temperatures of mass transfer line and injector were set at 220 and 300 °C, respectively. The oven temperature was initially set at 50 to 150 °C at a rate of 3 °C/min, then held for 10 min and finally raised to 300 °C at a rate of 10 °C/min. The crude samples was diluted with dichloromethane solvent (1:100, v/v) and filtered. The clean plant crude extracts (1 µL) were taken in a Hamilton syringe and injected into injector with split mode. The split ratio was 1:120. The percentage of chemical constituents in the plant crude was expressed as a percentage by peak area. Based on the retention time, the chemical constituents were identified and characterized in various crude extracts. The mass chromatograms of the crude extracts were matched with standards existing computer library (Mainlab, Replib and Tutorial data of GC-MS systems)[9-17].

3. Results

3.1. Analysis of different crude extracts

Ten different chemical constituents were identified in hexane crude extract from the leaves of *D. metel*, which are representing 5.54% of the total crude extract. The identified chemical constituents are listed in Table 1 according to their retention time. The different chemical constituents and chromatogram of hexane crude extract are showed in Figure 1 and Table 1. The chemical constituents are hexanoic acid (1.80%), heptanoic acid (0.31%), octanoic acid (0.08%), nonoic acid (0.21%), nonoic acid, z-oxo-ME (0.10%), eugenol (0.29%), neophytadiene (0.64%), hydrocarbon (1.59%), and vitamin E (0.52%).

Twenty five different chemical constituents were identified in ethyl acetate crude extract by GC-MS. The identified different chemical constituents represented 79.66% of the total extract. The list of chemical constituents and chromatogram were present in ethyl acetate crude extract (Figure 2 and Table 1), and the chemical constituents are as follows: pentanoic acid (0.93%), dihydroxy acetone (0.11%), hexanoic acid (18.40%), heptanoic acid (6.04%), nonanoic acid-2-oxo-mo (1.50%), octanoic acid (2.54%), glyco cyanidine (0.04%), nonanoic acid-ME (0.11%), nonanoic acid (6.69%), propanedioic acid (0.09%), decanoic acid (0.26%), β-endemol (0.47%), neophytadiene (12.02%), (2E)-3.7.11.15-tetramethyl-2-hexadecane (0.85%), hexadecanoic acid (1.14%), phytol (2.44%), α-linolenic acid-ME(z,z,z) (1.77%), squalene




Figure 1. Chemical constituents of hexane crude extract of *D. Metel*.




Figure 2. Chemical constituents of ethyl acetate crude extract of *D. metel*.




Figure 3. Chemical constituents of chloroform crude extract of *D. metel*.

(0.97%), hydrocarbon (6.70%), vitamin E (2a-tocophynol) (10.65%), and campesterol (5.94%). The chloroform crude extract from leaf samples of *D. metel* was analyzed and identified by using GC-MS, and the extract contained twenty one different chemical constituents

representing 87.34% of the total extract. The chemical constituents that were characterized in chloroform crude extract (Figure 3 and Table 1) are as hexanoic acid (39.67%), heptanoic acid (6.97%), hyoscyamine (10.88%), neophytadiene (6.61%) and nonoic acid (3.89%).




Figure 4. Chemical constituents of methanol crude extract of *D. metel*.

Table 1

Chemical constituents of different crude extracts from the leave of *D. metel*.

Compound Name	Retention time (min)	MW	Peak area (%)
Hexane crude extract	Hexanoic acid	6.64	1.80
	Heptanoic acid	9.78	0.31
	Octanoic acid	13.53	0.08
	Nonoic acid	17.58	0.21
	Nonoic acid, z-oxo-ME	20.84	0.10
	Eugenol	21.16	0.29
	Neophytadiene	38.89	0.64
	Hydrocarbon	70.55	0.41
	Hydrocarbon	71.34	1.18
	Vitamin E	72.09	0.52
	Chloroform crude extract	Pentanoic acid	4.19
Hexanoic acid		6.88	39.67
Heptanoic acid		9.95	6.97
Octanoic acid		13.63	1.78
Nonoic acid		19.37	3.89
2-methyl-4-vinylphenol		20.85	0.46
Nonoic acid, 2-oxo-ME		24.84	0.69
Tridecane, 4-8-dimethyl		25.85	1.19
Levogluconan		26.35	1.75
Neophytadiene		39.72	6.61
5-10-etoxy-2,3,7,8-tetrahydro-1H,6H,dipyrrolo(1,2-a:1:2-d)pyrazine		46.42	0.34
Anhydroatropine		50.74	1.01
Hyoscyamine		65.38	10.88
Squalene		67.09	0.36
Hydrocarbon		68.68	0.41
Hydrocarbon		70.55	0.72
Hydrocarbon		71.33	0.83
D- -tocophenol		71.33	2.63
Hydrocarbon		72.08	2.09
Hydrocarbon	72.83	1.74	
Hydrocarbon	73.37	0.75	
Methanol crude extract	Pentanoic acid	4.15	1.08
	Hexanoic acid-ME	5.18	0.52
	Octanol	5.68	0.66
	Hexanoic acid	6.70	27.93
	Heptanoic acid	9.86	6.11
	Pyranone	12.34	0.48
	Octonoic acid	13.53	2.20
	Nonoic acid	17.46	4.57

MW: Molecular weight.

Table 1, continued

Chemical constituents of different crude extracts from the leave of *D. metel*.

Compound Name	Retention time (min)	MW	Peak area (%)	
4-vinyl guaiacol	19.36	150	2.14	
Levogluconan	25.84	162	0.26	
Neophytadiene	38.89	278	7.30	
Plamitic acid	40.84	256	1.91	
Phytol	47.44	296	1.09	
-linolenic acid(z,z,z)	48.14	292	2.31	
Hyoscyamine	50.66	289	1.49	
Squalene	65.55	410	0.69	
Vitamine E	72.02	430	6.21	
Ethyl acetate crude extract	Pentanoic acid	4.19	102	0.93
	Dihydroxy acetone	4.40	90	0.11
	Hexanoic acid	6.99	116	18.40
	Heptanoic acid	10.06	130	6.04
	Nonanoic acid-2-oxo-mo	12.66	99	1.50
	Octanoic acid	1.73	144	2.54
	Glycocyantidine	15.43	99	0.04
	Nonanoic acid-ME	15.63	172	0.11
	Nonanoic acid	17.81	158	6.69
	Propanedioic acid	18.44	146	0.09
	Decanoic acid	21.53	172	0.26
	β -endesmol	32.34	222	0.47
	Neophytadiene	38.73	278	12.02
	(2E)-3,7,11,15-tetramethyl-2-hexadecane	39.01	280	0.85
	Hexadecanoic acid	42.79	256	1.14
	Phytol	47.30	296	2.44
	-linolenic acid-ME(z,z,z)	48.10	292	1.77
Squalene	65.39	410	0.97	
Hydrocarbon	66.94	420	0.37	
Hydrocarbon	68.52	420	1.43	
Hydrocarbon	70.38	420	1.17	
Hydrocarbon	71.16	420	1.11	
Vitamin E(2 α -tocophenol)	71.93	430	10.65	
Hydrocarbon	72.66	434	2.62	
Campesterol	73.70	400	5.94	

MW: Molecular weight.

Finally, the methanol crude extract was analyzed and identified by using GC-MS, and it contained seventeen different chemical constituent representing 66.95% of the total extract from leaf samples of *D. metel*. The chemical constituents founded in

the crude extract (Figure 4 and Table 1) were pentanoic acid (1.08%), hexanoic acid-ME (0.52%), octanol (0.66%), hexanoic acid (27.93%), heptanoic acid (6.11%), pyranone (0.48%), octanoic acid (2.20%), nonoic acid (4.57%), 4-vinyl guaiacol (2.14%), levoglucosan (0.26%), neophytadiene (7.30%), plamitic acid (1.91%), phytol (1.09%), α -linolenic acid(z,z,z) (2.31%), hyoscyamine (1.49%), squalene (0.69%), and vitamin E (6.21%).

3.2. Antimicrobial activity

Antibacterial potential of the prepared crude extracts was evaluated against three bacteria strains and the inhibition of bacterial growth was measured. Different crude extracts at different concentrations exhibited antibacterial potential against *S. aureus*, *E. coli* and *P. aeruginosa*. The antimicrobial results are presented in Table 2. All crude extracts from the leaves showed small potential against one Gram-positive and two Gram-negative bacteria at different concentrations with zones of inhibition in a range of 0-13 mm. Methanol crude extract showed small antibacterial potential against *E. coli* at all concentrations. However, the chloroform extracts did not show any antibacterial potential against *S. aureus* and *P. aeruginosa* at all concentrations. All the crude extracts from leaves showed moderate potential against *E. coli* at all working concentrations. The hexane and ethyl acetate crude extracts also showed moderate potential against *P. aeruginosa* but the remaining crude extracts did not show any potential against this bacteria. The ethyl acetate crude extract showed moderate activity against *S. aureus* at the concentration of 2 and 1 mg/mL. Butanol also showed activity against *S. aureus* at the concentrations 2.00, 1.00 and 0.50 mg/mL. However, the hexane and chloroform crude extracts did not show any activity against *P. aeruginosa* at any concentrations.

Table 2

Antimicrobial activity of different leaves crude extracts of *D. metel* against *E. coli*, *P. aeruginosa* and *S. aureus*

Crude extract	Concentration (mg/mL)	<i>E. coli</i> ^a (mm)	<i>S. aureus</i> (mm)	<i>P. aeruginosa</i> (mm)
Hexane	2.00	9.00±0.15	nd	9.00±0.17
	1.00	7.00±0.29	nd	7.00±0.21
	0.50	6.00±0.48	nd	7.00±0.11
	0.25	6.00±0.13	nd	6.00±0.24
	Standard	33.00±0.20	nd	dn
Chloroform	2.00	13.00±0.43	nd	dn
	1.00	8.00±0.06	nd	dn
	0.50	7.00±0.12	nd	dn
	0.25	6.00±0.17	nd	dn
	Standard	32.00±0.13	nd	dn
Ethyl acetate	2.00	7.00±0.22	8.00±0.90	7.00±0.19
	1.00	7.00±0.16	6.00±0.45	6.50±0.27
	0.50	6.00±0.18	dn	6.00±0.13
	0.25	6.00±0.23	dn	6.00±0.65
	Standard	28.00±0.31	dn	6.00±0.33
Butanol	2.00	9.00±0.13	9.00±0.20	dn
	1.00	9.00±0.07	7.00±0.16	dn
	0.50	8.00±0.28	6.00±0.26	dn
	0.25	7.00±0.43	dn	dn
	Standard	30.00±0.12	dn	dn
Methanol	2.00	8.00±0.18	7.00±0.38	dn
	1.00	8.00±0.12	dn	dn
	0.50	6.00±0.20	dn	dn
	0.25	6.00±0.11	dn	dn
	Standard	32.00±0.21	28.00±0.12	dn

nd: Not detected. Standard: Amoxicillin.

^a: Values are represented as mean±SD of three experiments.

4. Discussion

Different types of bioactive compounds are present in foodstuffs of both plant and animal origin. Epidemiological studies show a positive relationship between dietary intake of whole grains, fruits, vegetables, fish and fermented milk products and health status. All of these foodstuffs contain known bioactive compounds such as dietary fibres, phytosterols, carotenoids, peptides, bioactive lipids, alkaloid, flavonoids and probiotics[2-4]. The whole plant of *D. metel* contains high concentration of different alkaloids which increased gradually with maturity of the plant[17-25]. The yield of extraction by using different polarities solvents from the leaf of *D. metel* is different. The highest percentage of recovery yield is methanol and the lowest is ethyl acetate. According to the chromatogram, hexane crude extracts contain some bioactive compounds such as octanoic acid, eugenol and vitamin E which are widely used for preparation of different medicines[15-24]. The bioactive compounds, hexanoic acid (39.67%) and hyoscyamine (10.88%) isolated and identified from chloroform crude extracts have very strong antimicrobial property[14-18]. However, the activities of other compounds in the crude extract are still not known. The ethyl acetate and methanol crude extracts also contain lots of bioactive compounds which are traditionally and pharmaceutically used to prepare the medicine[22,23]. The detected compounds in the leaf crude extracts of *D. metel* may be responsible for the antioxidant and antibacterial potential. Several reports showed strong potential activities of the plant like antioxidant, anti-inflammatory, antimicrobial, antiangiogenic, anticancer and anti-allergic activities[21-23]. The antimicrobial potential of the plant crude extracts was measured by agar disc method against the mentioned three bacterial strains. The results obtained in this study are almost similar to antioxidant and antimicrobial activity of *D. metel* crude extracts reported by other authors[11,13,14].

No previous studies reported leaves extracts of *D. metel* grown in the Sultanate of Oman. In this study, a good number of bioactive compounds were isolated and identified from different polarities crude extracts. The methanol extract and its fractions possess moderate antimicrobial potential, and this may be due to the presence of alkaloids, normal hydrocarbons, cyclic aromatic hydrocarbons, organic acids, terpenoids and vitamin.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgements

The authors are grateful to Prof. Dr. Nafsiah Binti Shamsudin, Dean, College of Pharmacy and Nursing, University of Nizwa, Sultanate of Oman for the continuous encouragement during the work and providing all laboratory facilities. Authors also thank to Khaloud Ali Said Al-Alawi and Ahlam Rashed Alabri, Lab Technicians, Natural Product Lab, University of Nizwa for their continuous help during the experiment. The authors wish to express sincere gratitude to the Central Instrument Laboratory, College of Agriculture and Marine Sciences, Sultan Qaboos University, Sultanate of Oman where the test was confirmed (Grant No. 507/SOP/OB/1/2013).

Comments

Background

D. metel is the most important herbal plant used worldwide

for the treatment of different diseases. *D. metel* is a flowering plant belonging to family *Solanaceae*. Traditionally, it is also used for the treatment of catarrh, epilepsy, insanity, hysteria, rheumatic pains, hemorrhoids, painful menstruation, skin-ulcers and wounds. Currently, no work has been done on this species by the researcher.

Research frontiers

The aim of this study is to prepare various crude extracts using solvents with different polarities for qualitative evaluation of their chemical constituents by GC-MS, and study the antimicrobial potential of these crude extracts.

Related reports

Several report available showed strong potential activities of the plant like antioxidant, anti-inflammatory, antimicrobial, antiangiogenic, anticancer and anti-allergic properties.

Innovations and breakthroughs

The experimental data generated in this study is new information and data will be useful to the scientific community.

Applications

D. metel is used worldwide as a medicine. According to the present paper, there are so many bioactive compounds that can be used to prepare medicine.

Peer review

The present study on biochemical screening of various leaves crude extracts of *D. metel* is giving the valuable brief and scientific information about this plant.

References

- [1] Afsharypour S, Mostajeran A, Mokhtary R. Variation of scopolamine and atropine in different parts of *Datura metel* during development. *Planta Med* 1995; **61**: 383-384.
- [2] Akerele O. Summary of World Health Organization guidelines for the assessment of herbal medicines. *Herbalgram* 1993; **28**: 13-16.
- [3] Ali M, Shuaib M. Characterization of the chemical constituents of *Datura metel* Linn. *Indian J Pharm Sci* 1996; **58**(6): 243-245.
- [4] Hossain MA, Shah MD, Gnanaraj C, Iqbal M. *In vitro* total phenolics, flavonoids contents and antioxidant activity of essential oil, various organic extracts from the leaves of tropical medicinal plant *Tetragium* from Sabah. *Asian Pac J Trop Med* 2011; **4**: 717-721.
- [5] Hossain MA, Shah MD, Sang SV, Sakari M. Chemical composition and antibacterial properties of the essential oils and crude extracts of *Merremia borneensis*. *J King Saud Univ Sci* 2012; **24**: 243-249.
- [6] Anyasor GN, Ogunwenmo KO, Oyelana OA, Akpofunre BE. Phytochemical constituents and antioxidant activities of aqueous and methanol stem extracts of *Costus afer* Ker Gawl. (Costaceae). *Afr J Biotechnol* 2010; **9**(31): 4880-4884.
- [7] Ayoola GA, Coker HAB, Adesegun SA, Adepoju-Bello AA, Obaweya K, Ezennia EC, et al. Phytochemical screening and antioxidant activities of some selected medicinal plants used for malaria therapy in Southwestern Nigeria. *Trop J Pharm Res* 2008; **7**(3): 1019-1024.
- [8] Barile E, Bonanomi G, Antignani V, Zolfaghari B, Sajjadi SE, Scala F, et al. Saponins from *Allium minutiflorum* with antifungal activity. *Phytochemistry* 2007; **68**: 596-603.
- [9] Bhattacharjee I, Chatterjee SK, Chandra G. Isolation and identification of antibacterial components in seed extracts of *Argemone mexicana* L. (Papaveraceae). *Asian Pac J Trop Med* 2010; **3**(7): 547-551.
- [10] Bhimba BV, Meenupriya J, Joel EL, Naveena DE, Kumar S, Thangaraj M. Antibacterial activity and characterization of secondary metabolites isolated from mangrove plant *Avicennia officinalis*. *Asian Pac J Trop Med* 2010; **3**(7): 544-546.
- [11] Dabur R, Ali M, Singh H, Gupta J, Sharma GL. A novel antifungal pyrrole derivative from *Datura metel* leaves. *Pharmazie* 2004; **59**: 568-570.
- [12] Darabpour E, Motamedi H, Nejad SMS. Antimicrobial properties of *Teucrium polium* against some clinical pathogens. *Asian Pac J Trop Med* 2010; **3**(2): 124-127.
- [13] Anju D, Ratan L. Phytochemical and pharmacological status of *Datura fastuosa* Linn. *Int J Res Ayurveda Pharm* 2011; **2**(1): 145-150.
- [14] Okwu DE, Lgara EC. Isolation, characterization and antibacterial activity of alkaloid from *Datura metel* Linn leaves. *Afr J Pharm Pharmacol* 2009; **3**(5): 277-281.
- [15] Dorman HJ, Deans SG. Antimicrobial agents from plants: antibacterial activity of plant volatile oils. *J Appl Microbiol* 2000; **88**: 308-316.
- [16] Duke JA, Ayensu ES. *Medicinal plants of China*. Michigan: Reference Publ., Inc.; 1985, p. 90-91.
- [17] Hossain MA, Nagooru MR. Biochemical profiling and total flavonoids contents of leaves crude extract of endemic medicinal plant *Corydalis terminalis* L. Kunth. *Pharmacogn J* 2011; **3**(24): 25-30.
- [18] Hossain MA, Shah MD, Sakari M. Gas chromatography-mass spectrometry analysis of various organic extracts of *Merremia borneensis* from Sabah. *Asian Pac J Trop Med* 2011; **4**: 637-641.
- [19] Irudayaraj V, Janaky M, Johnson M, Selvan N. Preliminary phytochemical and antimicrobial studies on a spike-moss *Selaginella inaequalifolia* (hook. & grev.) spring. *Asian Pac J Trop Med* 2010; **3**(12): 957-960.
- [20] Igbinosa OO, Igbinosa EO, Aiyegoro OA. Antimicrobial activity and phytochemical screening of stem bark extracts from *Jatropha curcas* (Linn). *Afr J Pharm Pharmacol* 2009; **3**(2): 58-62.
- [21] Jarrar N, Abu-Hijleh A, Adwan K. Antibacterial activity of *Rosmarinus officinalis* L. alone and in combination with cefuroxime against methicillin-resistant *Staphylococcus aureus*. *Asian Pac J Trop Med* 2010; **3**(2): 121-123.
- [22] Jeeshna MV, Manorama S, Paulsamy S. Antimicrobial properties of the medicinal shrub, *Glycosmis pentaphylla*. *J Basic Appl Biol* 2009; **3**: 25-27.
- [23] Hashmi LS, Hossain MA, Weli AM, Al-Riyami Q, Al-Sabahi JN. Gas chromatography-mass spectrometry analysis of different organic crude extracts from the local medicinal plant of *Thymus vulgaris* L. *Asian Pac J Trop Biomed* 2013; **3**(1): 69-73.
- [24] Manandhar NP. *Plants and people of Nepal*. Portland: Timber Press; 2002.
- [25] Mann J. *Murder, magic and medicine*. Oxford: Oxford University Press; 2000.