

# Journal of Coastal Life Medicine

journal homepage: [www.jclmm.com](http://www.jclmm.com)



Document heading doi: 10.12980/JCLM.3.2015JCLM-2014-0083

©2015 by the Journal of Coastal Life Medicine. All rights reserved.

## Anaphylactoid reaction to biting midges bite

Sora Yasri<sup>1\*</sup>, Viroj Wiwanitkit<sup>2</sup>

<sup>1</sup>KMT Primary Care Center, Bangkok, Thailand

<sup>2</sup>Hainan Medical University, Haikou, China

### To the editor

Sir, biting midges is a common insect that can be seen in many beach areas in the tropical countries. Biting midges bite can cause health problem and this can be fatal. Here, the author reports a case of anaphylactoid reaction to biting midges bite. The patient was a 30 years' old male patient presenting with the problem of swelling right arm, palpitation and difficulty in respiration. This patient gave the history of biting midges bite at his right ring finger during having lunch at a restaurant at a beach. Within 5 minutes after getting bite, the patient felt serious pain and his hand and arm became swelling. His mother brought him to the physician and the patient started developing palpitation and respiratory difficulty during going to see the physician (about 10 min). The patient was diagnosed to have anaphylactoid reaction and the steroid therapy was given and the patient improved within 4 h. This is an interesting case of allergic reaction to biting midges bite. The saliva of biting midges contains several proteins which can cause the health problem[1]. Lehiy and Drolet noted for the understanding of roles of those proteins in "the possibly resulting disease pathogenesis[1]."

Chen *et al.* noted that "an immediate reaction to midge bites is immunoglobulin E-mediated. IFN-gamma, IL-6 and TNF-alpha are involved in delayed reactions to midge bites[2]." Of interesting, biting midges are abundant in many beaches in tropical zone and

there is no systematic control of those problematic insects[3]. To diagnose the disease, specific immunoglobulin E antibodies can be useful[4]. However, since the diagnostic tool is usually not available and the history of biting midges bite can be sufficient for diagnosis, the diagnosis is usually confirmed by clinical history as in the present case.

### Conflict of interest statement

We declare that we have no conflict of interest.

### References

- [1] Lehiy CJ, Drolet BS. The salivary secretome of the biting midge, *Culicoides sonorensis*. *PeerJ* 2014; **2**: e426.
- [2] Chen YH, Lee MF, Tsai JJ, Wu HJ, Hwang GY. Specific IgE and IgG responses and cytokine profile in subjects with allergic reactions to biting midge *Forcipomyia taiwana*. *Int Arch Allergy Immunol* 2009; **150**(1): 66-74.
- [3] Takoka H, Salazar NP, Cabrera BD. Observation on biting midge infestations on Mamburao Beach, Mindoro, Philippines. *Southeast Asian J Trop Med Public Health* 1976; **7**(4): 631-633.
- [4] Seldén AI, Calo A, Mølleby G, Hultgren O. Chironomid midge sensitization in sewage workers: case study. *Med Vet Entomol* 2013; **27**(3): 346-348.

\*Corresponding author: Sora Yasri, KMT Primary Care Center, Bangkok, Thailand.

E-mail: [sorayasri@outlook.co.th](mailto:sorayasri@outlook.co.th)

Article history:

Received 2 Jul 2014

Received in revised form 9 Jul, 2nd revised form 20 Jul, 3rd revised form 28 Jul 2014

Accepted 2 Oct 2014

Available online 2 Mar 2015